## antibodies .- online.com





## anti-IL1RAP antibody (AA 251-350) (Biotin)



O	:	
	rview	

Background:

Quantity:	100 μL
Target:	IL1RAP
Binding Specificity:	AA 251-350
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IL1RAP antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human IL1RAP/IL1R3
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat
Purification:	Purified by Protein A.
Target Details	
Target:	IL1RAP
Alternative Name:	IL1RAP (IL1RAP Products)

Synonyms: IL1R3, C3orf13, IL-1RAcP, Interleukin-1 receptor accessory protein, IL-1 receptor

accessory protein,	Interleukin-1	receptor 3	3, IL-1R-3,	IL-1R3, IL1RAP

Background: Coreceptor for IL1RL2 in the IL-36 signaling system (By similarity). Coreceptor with IL1R1 in the IL-1 signaling system. Associates with IL1R1 bound to IL1B to form the high affinity interleukin-1 receptor complex which mediates interleukin-1-dependent activation of NF-kappa-B and other pathways. Signaling involves the recruitment of adapter molecules such as TOLLIP, MYD88, and IRAK1 or IRAK2 via the respective TIR domains of the receptor/coreceptor subunits. Recruits TOLLIP to the signaling complex. Does not bind to interleukin-1 alone, binding of IL1RN to IL1R1, prevents its association with IL1R1 to form a signaling complex. The cellular response is modulated through a non-signaling association with the membrane IL1R2 decoy receptor. Secreted forms (isoforms 2 and 3) associate with secreted ligand-bound IL1R2 and increase the affinity of secreted IL1R2 for IL1B, this complex formation may be the dominant mechanism for neutralization of IL1B by secreted/soluble receptors.

Gene ID: 3556

UniProt: Q9NPH3

Pathways: NF-kappaB Signaling, Growth Factor Binding

## **Application Details**

Application Notes: IHC-P 1:200-400

Restrictions: For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months