

[Go to Product page](#)

Datasheet for ABIN2177394

anti-LPCAT2 antibody (AA 131-230) (Biotin)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | LPCAT2 |
| Binding Specificity: | AA 131-230 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This LPCAT2 antibody is conjugated to Biotin |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human LPCAT2 |
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse, Rat |
| Predicted Reactivity: | Cow, Sheep, Pig, Horse, Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|--|
| Target: | LPCAT2 |
| Alternative Name: | LPCAT2 (LPCAT2 Products) |

Target Details

| | |
|-------------|--|
| Background: | <p>Synonyms: AYTL1, LysoPAFAT, Lysophosphatidylcholine acyltransferase 2, LPC acyltransferase 2, LPCAT-2, LysoPC acyltransferase 2, 1-acylglycerol-3-phosphate O-acyltransferase 11, 1-AGP acyltransferase 11, 1-AGPAT 11, 1-acylglycerophosphocholine O-acyltransferase, 1-alkylglycerophosphocholine O-acetyltransferase, Acetyl-CoA:lyso-platelet-activating factor acetyltransferase, Acetyl-CoA:lyso-PAF acetyltransferase, Lyso-PAF acetyltransferase, Acyltransferase-like 1, Lysophosphatidic acid acyltransferase alpha, LPAAT-alpha, LPCAT2, AGPAT11</p> <p>Background: Possesses both acyltransferase and acetyltransferase activities. Activity is calcium-dependent. Involved in platelet-activating factor (PAF) biosynthesis by catalyzing the conversion of the PAF precursor, 1-O-alkyl-sn-glycero-3-phosphocholine (lyso-PAF) into 1-O-alkyl-2-acetyl-sn-glycero-3-phosphocholine (PAF). Also converts lyso-PAF to 1-O-alkyl-2-acyl-sn-glycero-3-phosphocholine (PC), a major component of cell membranes and a PAF precursor. Under resting conditions, acyltransferase activity is preferred. Upon acute inflammatory stimulus, acetyltransferase activity is enhanced and PAF synthesis increases. Also catalyzes the conversion of 1-acyl-sn-glycero-3-phosphocholine to 1,2-diacyl-sn-glycero-3-phosphocholine.</p> |
|-------------|--|

| | |
|----------|-------|
| Gene ID: | 54947 |
|----------|-------|

| | |
|----------|------------------------|
| UniProt: | Q7L5N7 |
|----------|------------------------|

Application Details

| | |
|--------------------|---|
| Application Notes: | WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500 |
| Restrictions: | For Research Use only |

Handling

| | |
|--------------------|---|
| Format: | Liquid |
| Concentration: | 1 µg/µL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |

Handling

| | |
|------------------|--------------------------------|
| | handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C for 12 months. |
| Expiry Date: | 12 months |