antibodies -online.com







Overview

Quantity:	100 μL
Target:	MAGI1
Binding Specificity:	AA 161-260
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MAGI1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MAGI1
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Dog,Cow,Sheep,Pig,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target: MAGI1

Target Details

Alternative Name:

MAGI1/IRSP58 (MAGI1 Products)

Background:

Synonyms: AIP 3, AIP3, Atrophin 1 interacting protein 3, atrophin-1 interacting protein 3, BAI associated protein 2, BAI1 associated protein 1, BAI1-associated protein 1, BAIAP 1, BAIAP1, BAP 1, BAP1, BAP2, Brain specific angiogenesis inhibitor 1 associated protein 2, Fas ligand associated factor 3, FLAF3, Gukmi1, Insulin receptor substrate p53, Insulin receptor substrate p53/p58, Insulin receptor substrate protein of 53 kDa, IRS 58, IRSP53, IRSp53/58, MAGI 1, MAGI-1, MAGI1c, Membrane associated quanylate kinase inverted 1, membrane associated guanylate kinase inverted-1, membrane associated guanylate kinase WW and PDZ domain containing 1, Protein BAP2, TNRC 19, TNRC19, Trinucleotide repeat containing gene 19, Trinucleotide repeat containing gene 19 protein, trinucleotide repeat-containing gene 19, WW domain containing protein 3, WW domain-containing protein 3, WWP 3, WWP3. Background: The membrane-associated guanylate kinase (MAGUK) proteins are concentrated at the membrane-cytoskeletal interface where they facilitate the assembly of multiprotein complexes on the inner surface of the plasma membrane. Three protein-protein interaction modules characteristically define MAGUK related proteins: the PDZ domain, the SH3 domain and the guanylate kinase (GuK) domain. The closely related MAGUK proteins, MAGI-1, MAGI-2 and MAGI-3 (membrane associated guanylate kinase inverted-1 and 2), likewise contain the GuK domain and five PDZ domains, however, the SH3 domain is replaced with a WW domain. The transcripts of MAGI-1 are alternatively spliced to produce three distinct proteins having unique C-terminals. Two variants, MAGI-1a and MAGI-1b, are associated with the membrane and cytosolic fractions and are primarily expressed in the brain. The third isoform, MAGI-1c, encodes for a nuclear localization signal that localizes MAGI-1c to the nucleus, and it is primarily expressed in the liver and kidney. MAGI-2 and MAGI-3 are localized to the plasma membrane, and they contribute to protein scaffolding by associating with the protein phosphatase PTEN.

Application Details

Application Notes: WB 1:300-5000

ELISA 1:500-1000

IHC-P 1:200-400

IHC-F 1:100-500

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

ICC 1:100-500

Application Details

Restrictions:	For Research Use only
Handling	
Format:	Liquid

Concentration:1 μg/μLBuffer:0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.Storage:4 °C,-20 °C

Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Expiry Date: 12 months