antibodies.com

# Datasheet for ABIN2178249 anti-PCDHB10 antibody (AA 351-450) (Biotin)



Overview

| Quantity:            | 100 µL   |
|----------------------|--|
| Target:              | PCDHB10  |
| Binding Specificity: | AA 351-450   |
| Reactivity:          | Human, Rat   |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This PCDHB10 antibody is conjugated to Biotin  |
| Application:         | Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)),<br>Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

# Product Details

| Immunogen:            | KLH conjugated synthetic peptide derived from human PCDHB10 |
|-----------------------|---|
| Isotype:              | lgG   |
| Cross-Reactivity:     | Human, Rat  |
| Predicted Reactivity: | Mouse,Rabbit  |
| Purification:         | Purified by Protein A.                                      |
| Target Details        |   |

| Target:           | PCDHB10                    |
|-------------------|----------------------------|
| Alternative Name: | PCDHB10 (PCDHB10 Products) |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN2178249 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

### Target Details

#### Background:

Synonyms: OTTHUMP00000160026, PCDBA\_HUMAN, PCDH BETA10, PCDH-beta-10, PCDHB10, PCHB10,Protocadherin beta-10, Protocadherin beta-10. Background: Protocadherins are a large family of cadherin-like cell adhesion proteins that are involved in the establishment and maintenance of neuronal connections in the brain. There are three protocadherin gene clusters, designated a, b and g, all of which contain multiple tandemlyarranged genes. PCDHB10 (protocadherin b10), also known as PCHB10 or PCDH-b10, is an 800 amino acid protein that is one of 16 proteins in the protocadherin b cluster. Unlike the a and g gene clusters, whose genes are spliced to downstream constant-region exons during transcription, members of the b cluster (such as PCDHB10) do not use constant-region exons to produce mRNAs. As a result, each protocadherin b gene encodes the transmembrane, extracellular and short cytoplasmic domains of the protein. Localized to the cell membrane, PCDHB10 is a single-pass type I membrane protein that contains six cadherin domains.

## Application Details

| Application Notes: | WB 1:300-5000         |
|--------------------|-----------------------|
|                    | IHC-P 1:200-400       |
|                    | IHC-F 1:100-500       |
| Restrictions:      | For Research Use only |

### Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.        |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C for 12 months.  |
| Expiry Date:       | 12 months  |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN2178249 | 03/07/2024 | Copyright antibodies-online. All rights reserved.