# antibodies - online.com







# anti-PCDH10 antibody (AA 21-120) (Biotin)



$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL	
Target:	PCDH10	
Binding Specificity:	AA 21-120	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PCDH10 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human PCDH10	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human,Cow,Sheep,Pig	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	PCDH10	
Alternative Name:	Protocadherin 10 (PCDH10 Products)	

## **Target Details**

Background:	Synonyms: PCDH19, OL-PCDH, Protocadherin-1, PCDH1, KIAA14	
	Background: Protocadherins are a large family of cadherin-like cell adhesion proteins that are	
	involved in the establishment and maintenance of neuronal connections in the brain. There are	
	three protocadherin gene clusters, designated alpha, beta and gamma, all of which contain	
	multiple tandemly arranged genes. PCDH10 (protocadherin 10), also known as PCDH19 or OL-	
	PCDH, is a 1,040 amino acid single-pass type I membrane protein that contains six cadherin	
	domains and one transmembrane domain. Expressed at moderate levels in brain, ovary and	
	testis and present at lower levels in all other tissues, PCDH10 functions as a calcium-dependent	
	cell-adhesion protein that may function as a tumor suppressor. When underexpressed, PCDH10	
	is associated with the progression of various carcinomas, including gastric cancer.	
Gene ID:	57575	
UniProt:	Q9P2E7	
Application Details		
Application Notes:	WB 1:300-5000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C for 12 months.	
Expiry Date:	12 months	