# antibodies .- online.com





Go to Product page

#### Datasheet for ABIN2178735

## anti-PCDH10 antibody (AA 21-120) (Cy5)

## Overview 100 μL Quantity: Target: PCDH10 Binding Specificity: AA 21-120 Reactivity: Rat Host: Rabbit Clonality: Polyclonal This PCDH10 antibody is conjugated to Cy5 Conjugate: Application: Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)) **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human PCDH10
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Cow,Sheep,Pig
Purification:	Purified by Protein A.

#### **Target Details**

Target:	PCDH10
Alternative Name:	Protocadherin 10 (PCDH10 Products)

### **Target Details**

-	
Background:	Synonyms: PCDH19, OL-PCDH, Protocadherin-1, PCDH1, KIAA14
	Background: Protocadherins are a large family of cadherin-like cell adhesion proteins that are
	involved in the establishment and maintenance of neuronal connections in the brain. There are
	three protocadherin gene clusters, designated alpha, beta and gamma, all of which contain
	multiple tandemly arranged genes. PCDH10 (protocadherin 10), also known as PCDH19 or OL-
	PCDH, is a 1,040 amino acid single-pass type I membrane protein that contains six cadherin
	domains and one transmembrane domain. Expressed at moderate levels in brain, ovary and
	testis and present at lower levels in all other tissues, PCDH10 functions as a calcium-dependent
	cell-adhesion protein that may function as a tumor suppressor. When underexpressed, PCDH10
	is associated with the progression of various carcinomas, including gastric cancer.
Gene ID:	57575
UniProt:	Q9P2E7
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months