# antibodies -online.com





Datasheet for ABIN2179275

## anti-SLK antibody (pTyr530)



( )	11/	IN	/ie	A .
	/ // <del> </del>	۱ ات	/   (−	' \/\/

Quantity:	100 μL
Target:	SLK
Binding Specificity:	pTyr530
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SLK antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffinembedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human FYN around the phosphorylation site of Tyr530
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Pig,Chicken,Rabbit
Purification:	Purified by Protein A.

#### Target Details

|--|

#### **Target Details**

Alternative Name:	SLK (SLK Products)		
Background:	Synonyms: Fyn phospho Y530, p-Fyn phospho Y530, C syn protooncogene, FYN oncogene		
	related to SRC FGR YES_ MGC45350_ OKT3 induced calcium influx regulator_ P59 FYN, Protein		
	tyrosine kinase fyn, Proto oncogene tyrosine protein kinase fyn, Protooncogene Syn, SLK, Src		
	like kinase, Src yes related novel gene, Src/yes related novel gene, SYN, FYN, Tyrosine kinase		
	p59fyn T, Tyrosine kinase p59fynT.		
	Background: Fyn is a membrane-associated, non-receptor protein tyrosine kinase		
	approximately 59 kDa, which belongs to the Src family of cytoplasmic tyrosine kinases. Fyn is		
	very strongly similar to mouse Fyn, v-yes and c-src. Fyn is expressed predominately in tissues		
	of neuronal and hematopoietic origin. Neuronal Fyn and hematopoietic Fyn differ at the junction		
	of the SH2 and kinase domains due to tissue specific alternative splicing. Fyn has been shown		
	to be involved in B cell and T cell activation as well as keratinocyte differentiation. In T cells, Fyr		
	associates with the T-cell antigen receptor and Thy1. The unique N terminal domain of Fyn		
	interacts with the CD3 and eta chains of the TcR. Fyn can bind to other proteins (p82 and p116)		
	through its SH2 and SH3 domains, which may act as substrates or regulators of Fyn activity.		
	Fyn is highly expressed in brain suggesting that it may have a role in the sensory nervous		
	network and in myelination at early stages of CNS formation.		
Gene ID:	2534		
Application Details			
Application Notes:	ELISA 1:500-1000		
	IHC-P 1:200-400		
	IHC-F 1:100-500		
	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.		
Preservative:	ProClin		

### Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months