
# antibodies -online.com





## anti-TAB3 antibody (PE)



#### Overview

Quantity:	100 μL
Target:	TAB3
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TAB3 antibody is conjugated to PE
Application:	Western Blotting (WB)

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human TAB3
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	TAB3
Alternative Name:	Tab3 (TAB3 Products)
Background:	Synonyms: MAP3K7IP 3, Mitogen activated protein kinase kinase kinase 7 interacting protein 3,
	Mitogen-activated protein kinase kinase kinase 7-interacting protein 3, NAP1, NF kappa B

activating protein 1, NF-kappa-B-activating protein 1, NFkB activating protein 1, TAB-3, TAB3,
TAB3_HUMAN, TAK1 binding protein 3, TAK1-binding protein 3, TGF-beta-activated kinase 1
and MAP3K7-binding protein 3, TGF-beta-activated kinase 1-binding protein 3.
Background: The product of this gene functions in the NF-kappaB signal transduction pathway.
The encoded protein, and the similar and functionally redundant protein MAP3K7IP2/TAB2,
forms a ternary complex with the protein kinase MAP3K7/TAK1 and either TRAF2 or TRAF6 in
response to stimulation with the pro-inflammatory cytokines TNF or IL-1. Subsequent
MAP3K7/TAK1 kinase activity triggers a signaling cascade leading to activation of the NF-
kappaB transcription factor. The human genome contains a related pseudogene. Alternatively
spliced transcript variants have been described, but their biological validity has not been
determined.
257397

Gene ID: 25739

UniProt: Q8N5C8

Pathways: TLR Signaling, Fc-epsilon Receptor Signaling Pathway, Activation of Innate immune Response,

Toll-Like Receptors Cascades

## **Application Details**

Application Notes:	FCM: (1:20-100)
	Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.