.-online.com antibodies

## Datasheet for ABIN2180075 anti-Vinexin antibody (AA 401-500) (Alexa Fluor 647)



Overview

Quantity:	100 µL	
Target:	Vinexin (Sorbs3)	
Binding Specificity:	AA 401-500	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Vinexin antibody is conjugated to Alexa Fluor 647	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Vinexin	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse	
Purification:	Purified by Protein A.	
Target Details		
Target:	Vinexin (Sorbs3)	
Alternative Name:	Vinexin (Sorbs3 Products)	

Background: Synonyms: SCAM 1, SCAM1, SH3 containing adapter molecule 1, SH3 domain containing

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN2180075 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

	adapter molecule 1, SH3D4, Sorbin and SH3 domain containing 3, Sorbin and SH3 domain	
	containing protein 3, Sorbs3, Vinexin alpha, Vinexin, Vinexin beta (SH3 containing adaptor molecule 1).	
	Background: Vinexin is a 671 amino acid protein that is expressed as two isoforms, designated	
	Vinexin alpha and Vinexin beta. Localized to cell junctions in both the cytoplasm and the	
	cytoskeleton, Vinexin alpha functions to promote Actin stress fiber formation, playing an	
	important role in modification of the Actin cytoskeleton. Like Vinexin alpha, Vinexin beta is	
	localized to cell junctions in the cytoplasm, but is also found in the nucleus where it plays an	
	important role in cell spreading and in activation of the JNK pathway in response to EGF	
	stimulation. Although Vinexin alpha and Vinexin beta have different roles within the cell, both	
	proteins contain three SH3 domains in their carboxy terminus and are expressed in a variety of	
	tissues, including placenta, heart, liver, brain, pancreas and skeletal muscle. Together, Vinexin	
	alpha and Vinexin beta are involved in cell-cell adhesion, signal transduction and cytoskeletal	
	organization throughout the cell.	
Pathways:	MAPK Signaling	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 µg/µL	
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN2180075 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

1.1	(	1:
Н	land	ling
		3

Expiry Date:

12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN2180075 | 03/07/2024 | Copyright antibodies-online. All rights reserved.