

Datasheet for ABIN2180410  
**anti-ZNF740 antibody (AA 101-193)**

## 2 Images

[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	ZNF740
Binding Specificity:	AA 101-193
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF740 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from Human ZNF740
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Cow, Sheep, Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	ZNF740
---------	--------

## Target Details


Alternative Name:	ZNF740 ( <a href="#">ZNF740 Products</a> )
Background:	Synonyms: MGC61706, OriLyt TD-element-binding protein 7, TB7, Zfp740, Zinc finger protein 740, ZN740_HUMAN, ZNF740.  Background: May be involved in transcriptional regulation.

## Application Details

Application Notes:	WB 1:300-5000 ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 ICC 1:100-500
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months



**Immunohistochemistry (Paraffin-embedded Sections)**

**Image 1.** Paraformaldehyde-fixed, paraffin embedded human cervical carcinoma tissue, Antigen retrieval by boiling in sodium citrate buffer(pH6) for 15min, Block endogenous peroxidase by 3% hydrogen peroxide for 30 minutes, Blocking buffer (normal goat serum) at 37°C for 20min, Antibody incubation with Rabbit Anti-ZNF740 Polyclonal Antibody, Unconjugated at 1:500 overnight at 4°C, followed by a conjugated secondary and DAB staining



**Immunohistochemistry (Paraffin-embedded Sections)**

**Image 2.** ZNF740 lysates probed with Polyclonal Antibody, unconjugated at 1:300 overnight at 4°C followed by a conjugated secondary antibody at 1:10000 for 90 minutes at 37°C.