antibodies - online.com







VTCN1 Protein (AA 29-258) (His tag)

Images



Overview

| Quantity: | 100 μg |
|-------------------------------|--|
| Target: | VTCN1 |
| Protein Characteristics: | AA 29-258 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Biological Activity: | Active |
| Purification tag / Conjugate: | This VTCN1 protein is labelled with His tag. |


Product Details

| Sequence: | AA 29-258 |
|------------------|--|
| Characteristics: | This protein carries a polyhistidine tag at the C-terminus. The protein has a calculated MW of 26.1 kDa. The protein migrates as 43-60 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation. |
| Purity: | >95 % as determined by SDS-PAGE. |
| Sterility: | 0.22 μm filtered |
| Endotoxin Level: | Less than 1.0 EU per μg by the LAL method. |

Target Details


Target Details

| V-set domain-containing T-cell activation inhibitor 1 (VTCN1) is also known as Immune costimulatory protein B7-H4, Protein B7S1, T-cell costimulatory molecule B7x, B7H4, which belongs to the immunoglobulin superfamily and BTN/MOG family. VTCN1 contains two Ig-like V-type (immunoglobulin-like) domains. The expression of VTCN1 is up-regulated by IL6 and IL10 and is inhibited by GM-CSF and IL4 on antigen-presenting cells (APCs). VTCN1 / B7-H4 negatively regulates T-cell-mediated immune response by inhibiting T-cell activation, proliferation, cytokine production and development of cytotoxicity. VTCN1 involved in promoting epithelial cell transformation. 26.4 kDa NP_078902 |
|--|
| belongs to the immunoglobulin superfamily and BTN/MOG family. VTCN1 contains two Ig-like V-type (immunoglobulin-like) domains. The expression of VTCN1 is up-regulated by IL6 and IL10 and is inhibited by GM-CSF and IL4 on antigen-presenting cells (APCs). VTCN1 / B7-H4 negatively regulates T-cell-mediated immune response by inhibiting T-cell activation, proliferation, cytokine production and development of cytotoxicity. VTCN1 involved in promoting epithelial cell transformation. |
| IL10 and is inhibited by GM-CSF and IL4 on antigen-presenting cells (APCs). VTCN1 / B7-H4 negatively regulates T-cell-mediated immune response by inhibiting T-cell activation, proliferation, cytokine production and development of cytotoxicity. VTCN1 involved in promoting epithelial cell transformation. |
| IL10 and is inhibited by GM-CSF and IL4 on antigen-presenting cells (APCs). VTCN1 / B7-H4 negatively regulates T-cell-mediated immune response by inhibiting T-cell activation, proliferation, cytokine production and development of cytotoxicity. VTCN1 involved in promoting epithelial cell transformation. |
| negatively regulates T-cell-mediated immune response by inhibiting T-cell activation, proliferation, cytokine production and development of cytotoxicity. VTCN1 involved in promoting epithelial cell transformation. 26.4 kDa |
| proliferation, cytokine production and development of cytotoxicity. VTCN1 involved in promoting epithelial cell transformation. 26.4 kDa |
| promoting epithelial cell transformation. 26.4 kDa |
| 26.4 kDa |
| |
| NP_078902 |
| |
| |
| For Research Use only |
| |
| Lyophilized |
| PBS, pH 7.4 |
| Please avoid repeated freeze-thaw cycles. |
| -20 °C |
| |
| Р |



SDS-PAGE

Image 1. Human B7-H4, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.



Binding Studies

Image 2. Immobilized Human B7-H4, His Tag with a linear range of 0.4-6 ng/mL.