

Datasheet for ABIN2180774

CD3 epsilon Protein (CD3E) (AA 22-117) (His tag)

2 Images




Overview

Quantity:	100 μg
Target:	CD3 epsilon (CD3E)
Protein Characteristics:	AA 22-117
Origin:	Cynomolgus
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This CD3 epsilon protein is labelled with His tag.
Product Details	
Sequence:	AA 22-117
Characteristics:	This protein carries a polyhistidine tag at the C-terminus. The protein has a calculated MW of
	11.7 kDa. The protein migrates as 18-20 kDa under reducing (R) condition (SDS-PAGE) due to
	glycosylation.
Purity:	>90 % as determined by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1.0 EU per μg by the LAL method.
Target Details	
Target:	CD3 epsilon (CD3E)


Target Details

rarget Details	
Alternative Name:	CD3 epsilon (CD3E Products)
Background:	CD3e molecule, epsilon is also known as CD3E, is a T-cell surface single-pass type I membrane
	glycoprotein. CD3E contains 1 lg-like (immunoglobulin-like) domain and 1 ITAM domain. CD3E,
	together with CD3-gamma, CD3-delta and CD3-zeta, and the T-cell receptor alpha/beta and
	gamma/delta heterodimers, forms the T cell receptor-CD3 complex. This complex plays an
	important role in coupling antigen recognition to several intracellular signal-transduction
	pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in the
	same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-cell
	development. CD3E plays an essential role in T-cell development, and defects in CD3E gene
	cause severe immunodeficiency. CD3E gene has also been linked to a susceptibility to type I
	diabetes in women. CD3E has been shown to interact with TOP2B, CD3EAP and NCK2.
Molecular Weight:	12.2 kDa
UniProt:	Q95LI5
Pathways:	TCR Signaling, CXCR4-mediated Signaling Events, Ubiquitin Proteasome Pathway
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Buffer:	PBS, pH 7.4
Handling Advice:	Please avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), After
	reconstitution under sterile conditions for 1 month (4 °C-8 °C) or 3 months (-20 °C to -70 °C).



ELISA

Image 1. Immobilized Cynomolgus CD3 epsilon, His Tag (ABIN2180775,ABIN2180774) at 2 μ g/mL (100 μ L/well) can bind Bispecific BCMAxCD3 T cing Antibody with a linear range of 0.4-6 ng/mL (QC tested).



SDS-PAGE

Image 2.