

## Datasheet for ABIN2180778

# CD3 epsilon Protein (CD3E) (AA 23-126) (His tag)

2 Images




## Overview

Quantity:	100 μg
Target:	CD3 epsilon (CD3E)
Protein Characteristics:	AA 23-126
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This CD3 epsilon protein is labelled with His tag.
Product Details	
Sequence:	AA 23-126
Characteristics:	This protein carries a polyhistidine tag at the C-terminus. The protein has a calculated MW of
	13.2 kDa. The protein migrates as 18-20 kDa under reducing (R) condition (SDS-PAGE) due to
	glycosylation.
Purity:	glycosylation. >90 % as determined by SDS-PAGE.
Purity: Sterility:	
•	>90 % as determined by SDS-PAGE.
Sterility:	>90 % as determined by SDS-PAGE.  0.22 µm filtered
Sterility: Endotoxin Level:	>90 % as determined by SDS-PAGE.  0.22 µm filtered


## **Target Details**

glycoprotein. CD3E contains 1 Ig-like (immunoglobulin-like) domain and 1 ITAM domain. together with CD3-gamma, CD3-delta and CD3-zeta, and the T-cell receptor alpha/beta a gamma/delta heterodimers, forms the T cell receptor-CD3 complex. This complex plays important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located ir same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-ce development. CD3E plays an essential role in T-cell development, and defects in CD3E ge cause severe immunodeficiency. CD3E gene has also been linked to a susceptibility to ty diabetes in women. CD3E has been shown to interact with TOP2B, CD3EAP and NCK2.  Molecular Weight: 12.6 kDa  NCBI Accession: NP_000724  Pathways: TCR Signalling, CXCR4-mediated Signalling Events, Ubiquitin Proteasome Pathway  Application Details  Restrictions: For Research Use only  Handling  Format: Lyophillized  Buffer: PBS, pH 7.4  Handling Advice: Please avoid repeated freeze-thaw cycles.  Storage: -20 °C  Storage Comment: No activity loss was observed after storage at: In lyophillized state for 1 year (4 °C-8 °C), 4 °C-8 °	Target Details	
glycoprotein. CD3E contains 1 Ig-like (immunoglobulin-like) domain and 1 ITAM domain. together with CD3-gamma, CD3-delta and CD3-zeta, and the T-cell receptor alpha/beta a gamma/delta heterodimers, forms the T cell receptor-CD3 complex. This complex plays important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located ir same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-ce development. CD3E plays an essential role in T-cell development, and defects in CD3E go cause severe immunodeficiency. CD3E gene has also been linked to a susceptibility to ty diabetes in women. CD3E has been shown to interact with TOP2B, CD3EAP and NCK2.  Molecular Weight: 12.6 kDa  NCBI Accession: NP_000724  Pathways: TCR Signaling, CXCR4-mediated Signaling Events, Ubiquitin Proteasome Pathway  Application Details  Restrictions: For Research Use only  Handling  Format: Lyophilized  Buffer: PBS, pH 7.4  Handling Advice: Please avoid repeated freeze-thaw cycles.  Storage: -20 °C  Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), A	Alternative Name:	CD3 epsilon (CD3E Products)
together with CD3-gamma, CD3-delta and CD3-zeta, and the T-cell receptor alpha/beta a gamma/delta heterodimers, forms the T cell receptor-CD3 complex. This complex plays important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypetides are located in same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-cel development, and defects in CD3E ge cause severe immunodeficiency. CD3E gene has also been linked to a susceptibility to ty diabetes in women. CD3E has been shown to interact with TOP2B, CD3EAP and NCK2.  Molecular Weight: 12.6 kDa  NCBI Accession: NP_000724  Pathways: TCR Signaling, CXCR4-mediated Signaling Events, Ubiquitin Proteasome Pathway  Application Details  Restrictions: For Research Use only  Handling  Format: Lyophilized  Buffer: PBS, pH 7.4  Handling Advice: Please avoid repeated freeze-thaw cycles.  Storage: -20 °C  Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), 4 °C-8	Background:	CD3e molecule, epsilon is also known as CD3E, is a T-cell surface single-pass type I membrane
gamma/delta heterodimers, forms the T cell receptor-CD3 complex. This complex plays important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-cell development, and defects in CD3E glays an essential role in T-cell development, and defects in CD3E gene has also been linked to a susceptibility to ty diabetes in women. CD3E has been shown to interact with TOP2B, CD3EAP and NCK2.  Molecular Weight: 12.6 kDa  NCBI Accession: NP_000724  Pathways: TCR Signaling, CXCR4-mediated Signaling Events, Ubiquitin Proteasome Pathway  Application Details  Restrictions: For Research Use only  Handling  Format: Lyophilized  Buffer: PBS, pH 7.4  Handling Advice: Please avoid repeated freeze-thaw cycles.  Storage: -20 °C  Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), // Storage Comment:		glycoprotein. CD3E contains 1 Ig-like (immunoglobulin-like) domain and 1 ITAM domain. CD3E
important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-cel development. CD3E plays an essential role in T-cell development, and defects in CD3E ge cause severe immunodeficiency. CD3E gene has also been linked to a susceptibility to ty diabetes in women. CD3E has been shown to interact with TOP2B, CD3EAP and NCK2.  Molecular Weight: 12.6 kDa  NCBI Accession: NP_000724  Pathways: TCR Signaling, CXCR4-mediated Signaling Events, Ubiquitin Proteasome Pathway  Application Details  Restrictions: For Research Use only  Handling  Format: Lyophilized  Buffer: PBS, pH 7.4  Handling Advice: Please avoid repeated freeze-thaw cycles.  Storage: -20 °C  Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), 4 °		together with CD3-gamma, CD3-delta and CD3-zeta, and the T-cell receptor alpha/beta and
pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-ce development. CD3E plays an essential role in T-cell development, and defects in CD3E gene cause severe immunodeficiency. CD3E gene has also been linked to a susceptibility to ty diabetes in women. CD3E has been shown to interact with TOP2B, CD3EAP and NCK2.  Molecular Weight: 12.6 kDa  NCBI Accession: NP_000724  Pathways: TCR Signaling, CXCR4-mediated Signaling Events, Ubiquitin Proteasome Pathway  Application Details  Restrictions: For Research Use only  Handling  Format: Lyophilized  Buffer: PBS, pH 7.4  Handling Advice: Please avoid repeated freeze-thaw cycles.  Storage: -20 °C  Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), A		gamma/delta heterodimers, forms the T cell receptor-CD3 complex. This complex plays an
same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-ce development. CD3E plays an essential role in T-cell development, and defects in CD3E ge cause severe immunodeficiency. CD3E gene has also been linked to a susceptibility to ty diabetes in women. CD3E has been shown to interact with TOP2B, CD3EAP and NCK2.  Molecular Weight: 12.6 kDa  NCBI Accession: NP_000724  Pathways: TCR Signaling, CXCR4-mediated Signaling Events, Ubiquitin Proteasome Pathway  Application Details  Restrictions: For Research Use only  Handling  Format: Lyophilized  Buffer: PBS, pH 7.4  Handling Advice: Please avoid repeated freeze-thaw cycles.  Storage: -20 °C  Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), A		important role in coupling antigen recognition to several intracellular signal-transduction
development. CD3E plays an essential role in T-cell development, and defects in CD3E ge cause severe immunodeficiency. CD3E gene has also been linked to a susceptibility to ty diabetes in women. CD3E has been shown to interact with TOP2B, CD3EAP and NCK2.  Molecular Weight: 12.6 kDa  NCBI Accession: NP_000724  Pathways: TCR Signaling, CXCR4-mediated Signaling Events, Ubiquitin Proteasome Pathway  Application Details  Restrictions: For Research Use only  Handling  Format: Lyophilized  Buffer: PBS, pH 7.4  Handling Advice: Please avoid repeated freeze-thaw cycles.  Storage: -20 °C  Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), A		pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in the
cause severe immunodeficiency. CD3E gene has also been linked to a susceptibility to ty diabetes in women. CD3E has been shown to interact with TOP2B, CD3EAP and NCK2.  Molecular Weight: 12.6 kDa  NCBI Accession: NP_000724  Pathways: TCR Signaling, CXCR4-mediated Signaling Events, Ubiquitin Proteasome Pathway  Application Details  Restrictions: For Research Use only  Handling  Format: Lyophilized  Buffer: PBS, pH 7.4  Handling Advice: Please avoid repeated freeze-thaw cycles.  Storage: -20 °C  Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), A		same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-cell
diabetes in women. CD3E has been shown to interact with TOP2B, CD3EAP and NCK2.  Molecular Weight: 12.6 kDa  NCBI Accession: NP_000724  Pathways: TCR Signaling, CXCR4-mediated Signaling Events, Ubiquitin Proteasome Pathway  Application Details  Restrictions: For Research Use only  Handling  Format: Lyophilized  Buffer: PBS, pH 7.4  Handling Advice: Please avoid repeated freeze-thaw cycles.  Storage: -20 °C  Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), A		development. CD3E plays an essential role in T-cell development, and defects in CD3E gene
Molecular Weight: 12.6 kDa  NCBI Accession: NP_000724  Pathways: TCR Signaling, CXCR4-mediated Signaling Events, Ubiquitin Proteasome Pathway  Application Details  Restrictions: For Research Use only  Handling  Format: Lyophilized  Buffer: PBS, pH 7.4  Handling Advice: Please avoid repeated freeze-thaw cycles.  Storage: -20 °C  Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), A		cause severe immunodeficiency. CD3E gene has also been linked to a susceptibility to type I
NCBI Accession: NP_000724  Pathways: TCR Signaling, CXCR4-mediated Signaling Events, Ubiquitin Proteasome Pathway  Application Details  Restrictions: For Research Use only  Handling  Format: Lyophilized  Buffer: PBS, pH 7.4  Handling Advice: Please avoid repeated freeze-thaw cycles.  Storage: -20 °C  Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), A		diabetes in women. CD3E has been shown to interact with TOP2B, CD3EAP and NCK2.
Pathways: TCR Signaling, CXCR4-mediated Signaling Events, Ubiquitin Proteasome Pathway  Application Details  Restrictions: For Research Use only  Handling  Format: Lyophilized  Buffer: PBS, pH 7.4  Handling Advice: Please avoid repeated freeze-thaw cycles.  Storage: -20 °C  Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), A	Molecular Weight:	12.6 kDa
Application Details  Restrictions: For Research Use only  Handling  Format: Lyophilized  Buffer: PBS, pH 7.4  Handling Advice: Please avoid repeated freeze-thaw cycles.  Storage: -20 °C  Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), A	NCBI Accession:	NP_000724
Restrictions: For Research Use only  Handling  Format: Lyophilized  Buffer: PBS, pH 7.4  Handling Advice: Please avoid repeated freeze-thaw cycles.  Storage: -20 °C  Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), A	Pathways:	TCR Signaling, CXCR4-mediated Signaling Events, Ubiquitin Proteasome Pathway
Handling  Format: Lyophilized  Buffer: PBS, pH 7.4  Handling Advice: Please avoid repeated freeze-thaw cycles.  Storage: -20 °C  Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), A	Application Details	
Format:  Lyophilized  Buffer:  PBS, pH 7.4  Handling Advice:  Please avoid repeated freeze-thaw cycles.  Storage:  -20 °C  Storage Comment:  No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), A	Restrictions:	For Research Use only
Buffer: PBS, pH 7.4  Handling Advice: Please avoid repeated freeze-thaw cycles.  Storage: -20 °C  Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), A	Handling	
Handling Advice:  Please avoid repeated freeze-thaw cycles.  Storage:  -20 °C  Storage Comment:  No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), A	Format:	Lyophilized
Storage: -20 °C  Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), A	Buffer:	PBS, pH 7.4
Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), A	Handling Advice:	Please avoid repeated freeze-thaw cycles.
	Storage:	-20 °C
reconstitution under sterile conditions for 1 month (4 °C-8 °C) or 3 months (-20 °C to -70	Storage Comment:	No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C-8 °C), After
		reconstitution under sterile conditions for 1 month (4 $^{\circ}$ C-8 $^{\circ}$ C) or 3 months (-20 $^{\circ}$ C to -70 $^{\circ}$ C).



### **ELISA**

**Image 1.** Immobilized Mouse A CD3 (SP34-2) at  $2 \mu g/mL$  (100  $\mu L/well$ ) can bind Human CD3 epsilon, His Tag (ABIN2180779,ABIN2180778) with a linear range of 0.4-6.25 ng/mL (QC tested).



#### **SDS-PAGE**

Image 2.