

Datasheet for ABIN2181607

**PD-1 Protein (AA 25-167) (Fc Tag)****3** Images**6** Publications[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	PD-1 (PDCD1)
Protein Characteristics:	AA 25-167
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This PD-1 protein is labelled with Fc Tag.

## Product Details

Sequence:	AA 25-167
Characteristics:	This protein carries a human IgG1 Fc tag at the C-terminus. The protein has a calculated MW of 42.6 kDa. The protein migrates as 50-66 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.
Purity:	>95 % as determined by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 0.1 EU per µg by the LAL method.

## Target Details

Target:	PD-1 (PDCD1)
---------	--------------

## Target Details

Alternative Name: PD-1 ([PDCD1 Products](#))

Background: Programmed cell death protein 1 (PD-1) is also known as CD279 and PDCD1, is a type I membrane protein and is a member of the extended CD28/CTLA-4 family of T cell regulators. PDCD1 is expressed on the surface of activated T cells, B cells, macrophages, myeloid cells and a subset of thymocytes. PD-1 has two ligands, PD-L1 and PD-L2, which are members of the B7 family. PD-L1 is expressed on almost all murine tumor cell lines, including PA1 myeloma, P815 mastocytoma, and B16 melanoma upon treatment with IFN- $\gamma$ . PD-L2 expression is more restricted and is expressed mainly by DCs and a few tumor lines. PD1 inhibits the T-cell proliferation and production of related cytokines including IL-1, IL-4, IL-10 and IFN- $\gamma$  by suppressing the activation and transduction of PI3K/AKT pathway. In addition, coligation of PD1 inhibits BCR-mediating signal by dephosphorylating key signal transducer. In vitro, treatment of anti-CD3 stimulated T cells with PD-L1-Ig results in reduced T cell proliferation and IFN- $\gamma$  secretion. Monoclonal antibodies targeting PD-1 that boost the immune system are being developed for the treatment of cancer.

Molecular Weight: 42.1 kDa

NCBI Accession: [NP\\_005009](#)

Pathways: [Cancer Immune Checkpoints](#)

## Application Details

Restrictions: For Research Use only

## Handling

Format: Lyophilized

Buffer: PBS, pH 7.4

Handling Advice: Please avoid repeated freeze-thaw cycles.

Storage: -20 °C

Storage Comment: No activity loss was observed after storage at: In lyophilized state for 1 year (4 °C), After reconstitution under sterile conditions for 3 months (-70 °C).


## Publications

Product cited in: Mettler Izquierdo, Varela, Park, Collarini, Lu, Pramanick, Rucker, Lopalco, Etches, Harriman: "High-efficiency antibody discovery achieved with multiplexed microscopy." in: **Microscopy**

(Oxford, England), Vol. 65, Issue 4, pp. 341-52, (2018) ([PubMed](#)).


There are more publications referencing this product on: [Product page](#)

Images




SDS-PAGE

**Image 1.** Human PD-1, Fc Tag (HPLC-verified) on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.



Binding Studies

**Image 2.** Immobilized Human PD-L1, His Tag with a linear range of 0.05-1 µg/mL.



Binding Studies

**Image 3.** Immobilized Human PD-L2, His Tag with a linear range of 0.02-0.5 µg/mL.