

# Datasheet for ABIN2191753

# anti-CD31 antibody





Go to Product page

$\sim$				
( )\	/e	r\ /	$\cap$	٨
()	<i>v</i> $\leftarrow$	ı vı	$\Box$	٧V

Our and the second	100	
Quantity:	100 μg	
Target:	CD31 (PECAM1)	
Reactivity:	Mouse	
Host:	Rat	
Clonality:	Monoclonal	
Conjugate:	This CD31 antibody is un-conjugated	
Application:	Flow Cytometry (FACS), Immunofluorescence (IF), Immunoprecipitation (IP)	
Product Details		
Clone:	MEC7-46	
Sterility:	0.2 μm filtered	
Target Details		
Target:	CD31 (PECAM1)	
Alternative Name:	Pecam-1 (PECAM1 Products)	
Background:	The monoclonal antibody MEC7.46 recognizes the mouse form of the platelet-endothelial adhesion molecule (PECAM)-1 (CD31). PECAM-1 is a member of the immunoglobulin superfamily. This heavily glycosylated protein is found in the entire vascular endothelium adult mice and functions in mediating cellular adhesion by heterophilic and homophilic mechanisms. PECAM-1 is detected within the lymphopoietic islands in the spleen of new (day 12) and in the bone marrow of adult mice. Capillary endothelial cells of adult mice along the control of the platelet-endothelia.	

express PECAM-1. The reactivity of the monoclonal antibody MEC7.46 is restricted to the isoform of the molecule that is selectively expressed by endothelial cells. The antibody precipitates a 130 kDa molecule present on the membrane of endothelial cells of all mouse blood vessels both in normal, inflamed and tumor tissues. The antigen is predominantly present at the lateral borders of endothelial cells as described for human PECAM-1. Staining of MEC7.46 can be seen on capillaries, veins, arteries and liver sinusoids. Aliases CD31, plateletendothelial cell adhesion molecule-1 Immunogen Mouse t.end.1 cells (polyoma middle T (PmT)-transformed EC)

Pathways:

Regulation of Actin Filament Polymerization

## **Application Details**

**Application Notes:** 

For immunohistochemistry and flow cytometry, dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. Positive Mouse t-end.1 cells control Negative L929 cells (Fibroblasts) control 1

Restrictions:

For Research Use only

#### Handling

Buffer:

PBS, containing 0.1 % bovine serum albumin and 0.02 % sodium azide.

Preservative:

Sodium azide

Precaution of Use:

This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage:

4°C

Storage Comment:

Product should be stored at 4 °C. Under recommended storage conditions, product is stable for at least one year. The exact expiry date is indicated on the label.

### **Publications**

Product cited in:

Xu, Zhang, Davison, Hu: "Circulating progenitor cells regenerate endothelium of vein graft atherosclerosis, which is diminished in ApoE-deficient mice." in: **Circulation research**, Vol. 93, Issue 8, pp. e76-86, (2003) (PubMed).