# antibodies -online.com





## anti-nectin-3 antibody

### **Publications**



( )	11	$\sim$	rv		۱ ۸
	1 \ /	┙	I \/	╙	1/1

Background:

Overview	
Quantity:	100 μg
Target:	nectin-3 (NECTIN3)
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This nectin-3 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunofluorescence (IF), Immunoassay (IA)
Product Details	
Clone:	103-A1
Isotype:	lgG2a
Cross-Reactivity (Details):	Cross reactivity: Human nectin-3 : No, Rat nectin-3 : No
Sterility:	0.2 μm filtered
Target Details	
Target:	nectin-3 (NECTIN3)
Alternative Name:	

The monoclonal antibody 103-A1 recognizes mouse nectin-3. Nectin-3 is a 83 kDa type I

transmembrane glycoprotein. Nectin, originally isolated as poliovirus receptor-related protein

(PRR), is a cell-cell adhesion molecule of the immunoglobulin supergene family. Nectins are

calcium- independent immunoglobulin-like cell-cell adhesion molecules consisting of four

members, nectin 1-4. Nectins homophilically and heterophilically trans-interact to form a variety of cell-cell junctions, including cadherin-based adherens junctions in epithelial cells and fibroblasts in culture, synaptic junctions in neurons, and Sertoli cell-spermatid junctions in testis, in cooperation with, or independently of, cadherins. Both nectin-2 and nectin-3 are ubiquitously expressed, whereas nectin-1 is abundantly expressed in brain. Nectin-2 and -3 are expressed in cells where cadherin is not expressed, such as blood cells and spermatids. All members of the nectin family have two or three splice variants. For nectin-3, three isoforms exist: nectin-3α, -3β and -3γ of which nectin-3α is the largest. Nectin-3, also known as PRR3, is a transmembrane protein that is predominantly expressed in testis and placental tissues as well in many cell lines. Nectin interacts in vivo with both long and short isoforms of afadin, an actin binding protein, at cadherin-based cell-cell adherence junctions in various tissues and cell lines. Furthermore, the ectodomains of nectin-3 and CD155 (Poliovirus Receptor) have shown strong affinity to each other. Injection of antibody 103-A1 into lumen of seminiferous tubules leads to disruption of the actin filaments in Sertoli cells at the Sertoli-maturing spermatid ectoplasmic specialization and exfoliation of maturing spermatids form the seminiferous epithelium. Aliases PPR3, poliovirus receptor-related protein 3, PRR3, PVRR3, CD113 Immunogen Recombinant mouse nectin-3 extracellular domain (50-400 aa) fused to IgG Fc

Pathways:

Cell-Cell Junction Organization

#### **Application Details**

**Application Notes:** 

For immunohistochemistry and flow cytometry, dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. Positive L-cells (A cultured line of C3H mouse fibroblasts), control Negative Cells/tissues derived from nectin-3 knockout mouse (Ref 5) control

Restrictions:

For Research Use only

#### Handling

Buffer:	PBS, containing 0.1 % bovine serum albumin and 0.02 % sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C	

#### Handling

#### Storage Comment:

Product should be stored at 4 °C. Under recommended storage conditions, product is stable for at least one year. The exact expiry date is indicated on the label.

#### **Publications**

#### Product cited in:

Feucht, Schneeberger, Hillebrand, Burkhardt, Weiss, Riethmüller, Land, Albert: "Capillary deposition of C4d complement fragment and early renal graft loss." in: **Kidney international**, Vol. 43, Issue 6, pp. 1333-8, (1993) (PubMed).

Zwirner, Felber, Herzog, Riethmüller, Feucht: "Classical pathway of complement activation in normal and diseased human glomeruli." in: **Kidney international**, Vol. 36, Issue 6, pp. 1069-77, (1990) (PubMed).