



Datasheet for ABIN2191806

anti-JAM3 antibody



[Go to Product page](#)

1 Publication

Overview

Quantity:	100 µg
Target:	JAM3
Reactivity:	Mouse
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This JAM3 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunoprecipitation (IP), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunoassay (IA)

Product Details

Clone:	CRAM-19 H36
Sterility:	0.2 µm filtered

Target Details

Target:	JAM3
Alternative Name:	Junctional Adhesion Molecule-C (JAM3 Products)
Background:	Junctional adhesion molecule-C (JAM-C) also known as JAM-2 is a 45 kD cell adhesion molecule (CAM). JAM-C is a transmembrane protein which is a member of the immunoglobulin superfamily found at intercellular junctions of endothelial cells. JAM-C belongs together with JAM-A (JAM or JAM-1) and JAM-B (VE-JAM or JAM-3) to a family of adhesion proteins with a V-C2 immunoglobulin domain organization. JAM plays an important role in tight junctions

Target Details

where it is involved in cell-to-cell adhesion through homophilic interaction. It codistributes with other tight junction components as ZO-1, 7H6 antigen, cingulin and occludin. JAM-C is potentially involved in the junctional sealing of the vascular endothelium, in particular of high endothelial venules (HEV). In adult murine tissue JAM-C expression is reported to be restricted to high endothelial venules of lymphoid organs, lymphoendothelial cells and endothelial cells in kidney. Monoclonal antibody CRAM-19 H36 also reacts with human JAM-C. In humans, JAM-C expression is not restricted to endothelial cells, but is also expressed on human lymphocytes.

Application Details

Application Notes: For immunohistology and flow cytometry dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.

Restrictions: For Research Use only

Handling

Buffer: PBS, containing 0.1 % bovine serum albumin and 0.02 % sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C

Storage Comment: Product should be stored at 4 °C. Under recommended storage conditions, product is stable for one year.

Expiry Date: 12 months

Publications

Product cited in: Feucht, Schneeberger, Hillebrand, Burkhardt, Weiss, Riethmüller, Land, Albert: "Capillary deposition of C4d complement fragment and early renal graft loss." in: **Kidney international**, Vol. 43, Issue 6, pp. 1333-8, (1993) ([PubMed](#)).

Zwirner, Felber, Herzog, Riethmüller, Feucht: "Classical pathway of complement activation in normal and diseased human glomeruli." in: **Kidney international**, Vol. 36, Issue 6, pp. 1069-77, (1990) ([PubMed](#)).

