.-online.com antibodies

Datasheet for ABIN2191899 anti-Cadherin 5 antibody (FITC)

2 Images

6 Publications



Overview

Quantity:	100 µg
Target:	Cadherin 5 (CDH5)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Cadherin 5 antibody is conjugated to FITC
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (IF), Functional Studies (Func), Immunoassay (IA)

Product Details

Clone:	BV9
Sterility:	0.2 µm filtered

Target Details

Target:	Cadherin 5 (CDH5)
Alternative Name:	Vascular Endothelial Cadherin (CDH5 Products)
Background:	The monoclonal antibody BV9 binds to the extracellular domain (EC3-EC4) of human VE-
	cadherin (vascular endothelial cadherin). Endothelial cells control the passage of plasma
	constituents and circulating cells from blood to the underlying tissues. VE-cadherin is of vital
	importance for the maintenance and control of endothelial cell contacts. Mechanisms that
	regulate VE-cadherin- mediated adhesion are important for the control of vascular permeability

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN2191899 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

and leukocyte extravasation. VE-cadherin regulates various cellular processes such as cell proliferation and apoptosis and modulates vascular endothelial growth factor receptor functions. Therefore, VE-cadherin is also essential during embryonic angiogenesis. The specialized function of VE-cadherin is lost or impaired in several pathological conditions including inflammation, sepsis, ischemia and diabetes - which leads to severe, and sometimes fatal, organ dysfunction. Furthermore, abnormal increase in vascular permeability is often observed in pathological conditions, such as tumor-induced angiogenesis, macular degeneration, allergy, and brain stroke. Endothelial permeability is regulated in part by the dynamic opening and closure of cell-cell adherent junctions. In vascular endothelium, adherent junctions are mainly composed of VE-cadherin, an adhesive receptor that is able to selfassociate at endothelial cell-cell contacts. VE-cadherin links endothelial cells together by homophilic interactions mediated by its extracellular part and associates intracellularly with the actin cytoskeleton via catenins. VE-cadherin belongs to the cadherin super- family of cell-cell adhesion molecules, which are encoded by more than 200 genes in the human 2+ genome. Classical cadherins are Ca -dependent, homophilic, cell to cell adhesion molecules expressed in nearly all cells within solid tissues. Cadherins form a core adhesion complex that consists of a cadherin dimer, binding through its extracellular region to another dimer of cadherins expressed in adjacent cells, while its intracellular region is anchored to the plasma membrane and linked to the cytoskeleton. The VE-cadherin extracellular domain consists of five cadherin-type repeats, called EC (extracellular cadherin) domains that are bound together by calcium ions in a rod-like structure. Aliases 7B4 antigen, Cadherin-5, CD144

Pathways:

Cell-Cell Junction Organization, Signaling Events mediated by VEGFR1 and VEGFR2

Application Details

Application Notes:	For immunohistochemistry, flow cytometry and Western blotting, dilutions to be used depend
	on detection system applied. It is recommended that users test the reagent and determine their
	own optimal dilutions. The typical starting working dilution is 1:50. For functional studies, in
	vitro dilutions have to be optimized in user's experimental setting. Positive HUVECs grown on
	coverslips control
Restrictions:	For Research Use only
Handling	
Buffer:	PBS, containing 1 % bovine serum albumin and 0.02 % sodium azide.
Preservative:	Sodium azide

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN2191899 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

Handling	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Product should be stored at 4 °C. Under recommended storage conditions, product is stable for at least one year. The exact expiry date is indicated on the label.
Publications	
Product cited in:	Nascimento, Sallé, Hoebeke, Argibay, Peineau: "cGMP-mediated inhibition of cardiac L-type Ca(2+) current by a monoclonal antibody against the M(2) ACh receptor." in: American journal of physiology. Cell physiology , Vol. 281, Issue 4, pp. C1251-8, (2001) (PubMed).
	Elies, Fu, Eftekhari, Wallukat, Schulze, Granier, Hjalmarson, Hoebeke: "Immunochemical and functional characterization of an agonist-like monoclonal antibody against the M2 acetylcholine receptor." in: European journal of biochemistry / FEBS , Vol. 251, Issue 3, pp. 659-66, (1998) (PubMed).

There are more publications referencing this product on: Product page

Images



Flow Cytometry

Image 1. Figure 1. Endothelial cells were fixed with 4% PAF (15 min, room temp), and then permeabilized with 0.5% TritionX-100 (3 min, room temperature). Cells were incubated with a final concentration of 10mg/ml BV9. Secondary detection was performed with anti-mouse Alexa-Fluor 647 and counterstained using DAPI.

Figure 2. HUVEC cells were stained with antibody BV9 in PBS/0.1% saponin for 1h at 4°C.(Black- no stain, Red- is otype control, Blue- HM2032–BV9)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/4 | Product datasheet for ABIN2191899 | 09/11/2023 | Copyright antibodies-online. All rights reserved.



Immunofluorescence

Image 2. Figure 1. Endothelial cells were fixed with 4% PAF (15 min, room temp), and then permeabilized with 0.5% TritionX-100 (3 min, room temperature). Cells were incubated with a final concentration of 10mg/ml BV9. Secondary detection was performed with anti-mouse Alexa-Fluor 647 and counterstained using DAPI.

Figure 2. HUVEC cells were stained with antibody BV9 in PBS/0.1% saponin for 1h at 4°C.(Black- no stain, Red- is otype control, Blue- HM2032–BV9)