# antibodies -online.com







## **Publications**



Go to Product page

_					
U	V	er	V	Ie	W

Quantity:	100 μg
Target:	MBP/MBL
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This MBP/MBL antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunohistochemistry (Frozen Sections) (IHC (fro)), Flow Cytometry (FACS), Immunoassay (IA)

#### **Product Details**

Clone:	3E7
Sterility:	0.2 μm filtered

### **Target Details**

Target:

Alternative Name:	MbI (MBP/MBL Products)
Background:	Mannose Binding Lectin (MBL) also called mannose- or mannan-binding protein (MBP) is a
	member of the group of collectins. MBL is an oligomeric lectin that recognizes carbohydrates
	as mannose and N- acetylglucosamine on pathogens. MBL contains a cysteine rich, a collagen
	like and a carbohydrate recognition domain. It forms a complex with C1r/C1s like serine
	proteases designated MASPs that proteolytically cleave C4_C2 and C3_MRL is able to activate

MBP/MBL

#### **Target Details**

the complement pathway independent of the classical and alternative complement activation pathways. The MBL-MASP pathway (better known as the lectin pathway) is antibody and C1q-independent. MBL exhibits complement-dependent antibacterial activity and acts directly as an opsonic and therefore plays an important role in innate immunity. MBL is synthesized by hepatocytes and has been isolated from the liver or serum of various vertebrate species.

#### **Application Details**

Δnn	lication	Notas.
App	lication	MOLES.

For Western blotting, flow cytometry and immunohistology dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.

Restrictions:

For Research Use only

#### Handling

Buffer:	PBS, containing 1 % bovine serum albumin and 0.02 % sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C	
Storage Comment:	Product should be stored at 4 °C. Under recommended storage conditions, product is stable for one year.	
Expiry Date:	12 months	

#### **Publications**

Product cited in:

Bax, Siersema, Haringsma, Kuipers, Vos, Van Dekken, Van Vliet, Kusters: "High-grade dysplasia in Barrett's esophagus is associated with increased expression of calgranulin A and B." in:

Scandinavian journal of gastroenterology, Vol. 42, Issue 8, pp. 902-10, (2007) (PubMed).