

Datasheet for ABIN2192082 anti-PTX3 antibody

5 Publications



## Overview

Quantity:	100 µg
Target:	PTX3
Reactivity:	Human
Host:	Rat
Clonality:	Monoclonal
Conjugate:	This PTX3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunoprecipitation (IP), Immunofluorescence (IF), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry
	(Frozen Sections) (IHC (tro)), Flow Cytometry (FACS), Immunoassay (IA)

## Product Details

Clone:	MNB4
Sterility:	0.2 µm filtered

## Target Details

Target:	PTX3
Alternative Name:	Pentraxin 3 (PTX3 Products)
Background:	The monoclonal antibody MNB4 (previously known as clone 20) recognizes human pentraxin 3
	(PTX3), belonging to the long pentraxin family. PTX3 is an acute-phase glycoprotein of ${\sim}45{ m kDa}$
	with glycosylation accounting for about 10 % of its molecular weight. PTX3 has a complex
	oligomeric structure with protomers linked to each other by disulfide bonds. PTX3 expression is

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN2192082 | 07/26/2024 | Copyright antibodies-online. All rights reserved. triggered by inflammatory cytokines, resulting in higher levels of circulating PTX3. Several cell types have been reported to produce PTX3, namely macrophages, endothelial cells, neutrophils and synoviocytes. PTX3 is involved in host defense against pathogen infection, in the regulation of the scavenger activity of macrophages and dendritic cells, and in modulation of complement activity by binding to C1q. Furthermore, PTX3 has been implicated in matrix deposition of cumulus cells. Moreover, PTX3 interacts with other biologically active molecules, causing their functional blockade. This has been demonstrated for fibroblast growth factor-2 (FGF-2), for which PTX3 acts as an inhibitor, leading to inhibition of angiogenesis. PTX3, like other pentraxins C-reactive protein (CRP) and serum amyloid P component (SAP), binds apoptotic cells and debris. PTX3 is useful as an early indicator of myocyte irreversible injury in ischemic cardiomyopathy. PTX3 is not only involved in inflammatory vessel diseases related to atherosclerosis, but also in pre-eclampsia and systemic small vessel ANCA-associated vasulitis, in which neutrophils are key players. The relationship between tissue damage and pentraxin generation is stringent in acute injuries. PTX3 tunes self-non-self discrimination and tissue repair due to the recognition of diverse ligands by PTX3 and through regulation of effector pathways. Aliases Pentraxin-related protein PTX3, Tumor necrosis factor-inducible gene 14 protein, TSG-14 Immunogen Human recombinant PTX3

## **Application Details**

Application Notes:	For immunohistology, flow cytometry and Western blotting, dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50
Restrictions:	For Research Use only
Handling	
Buffer:	PBS, containing 0.1 % bovine serum albumin and 0.02 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Product should be stored at 4 °C. Under recommended storage conditions, product is stable for one year.
Expiry Date:	12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN2192082 | 07/26/2024 | Copyright antibodies-online. All rights reserved. Product cited in:

Jaillon, Peri, Delneste, Frémaux, Doni, Moalli, Garlanda, Romani, Gascan, Bellocchio, Bozza, Cassatella, Jeannin, Mantovani: "The humoral pattern recognition receptor PTX3 is stored in neutrophil granules and localizes in extracellular traps." in: **The Journal of experimental medicine**, Vol. 204, Issue 4, pp. 793-804, (2007) (PubMed).

Inforzato, Peri, Doni, Garlanda, Mantovani, Bastone, Carpentieri, Amoresano, Pucci, Roos, Daha, Vincenti, Gallo, Carminati, De Santis, Salvatori: "Structure and function of the long pentraxin PTX3 glycosidic moiety: fine-tuning of the interaction with C1q and complement activation." in: **Biochemistry**, Vol. 45, Issue 38, pp. 11540-51, (2006) (PubMed).

Bussolati, Peri, Salvidio, Verzola, Mantovani, Camussi: "The long pentraxin PTX3 is synthesized in IgA glomerulonephritis and activates mesangial cells." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 170, Issue 3, pp. 1466-72, (2003) (PubMed).

Fazzini, Peri, Doni, DellAntonio, Dal Cin, Bozzolo, DAuria, Praderio, Ciboddo, Sabbadini, Manfredi, Mantovani, Querini: "PTX3 in small-vessel vasculitides: an independent indicator of disease activity produced at sites of inflammation." in: **Arthritis and rheumatism**, Vol. 44, Issue 12, pp. 2841-50, (2001) (PubMed).

Peri, Introna, Corradi, Iacuitti, Signorini, Avanzini, Pizzetti, Maggioni, Moccetti, Metra, Cas, Ghezzi, Sipe, Re, Olivetti, Mantovani, Latini: "PTX3, A prototypical long pentraxin, is an early indicator of acute myocardial infarction in humans." in: **Circulation**, Vol. 102, Issue 6, pp. 636-41 , (2000) (PubMed).