# antibodies .- online.com



Datasheet for ABIN2344856

# CytoSelect™ 96-well Cell Migration and Invasion Assay (8 µm), Fluorometric, Combo Kit



Go to Product page

# 6 F

## **Publications**

### Overview

Quantity:	2 x 96 tests
Reactivity:	Mammalian
Application:	Cellular Assay (CA)

#### **Product Details**

Brand:	CytoSelect™
Sample Type:	Serum, Cell Samples
Analytical Method:	Quantitative
Detection Method:	Fluorometric
Characteristics:	CytoSelect™ 96-well Cell Migration and Invasion Assay Kit utilizes a polycarbonate membrane
	plate (8 µm pore size) or basement membrane-coated inserts to assay the migratory or
	invasive properties of cells. The kit does not require you to prelabel the cells with Calcein AM or
	remove non-migratory or non-invasive cells (i.e. cotton swabbing). Any migratory or invasive
	cells are first dissociated from the membrane, then lysed and detected with CyQuant® GR Dye.
	The CytoSelect™ 96-well Cell Migration and Invasion Assay Kit provides a robust system for the
	quantitative determination of cell migration. Each assay contains sufficient reagents for the
	evaluation of 96 samples. The 8 µm pore size is optimal for epithelial and fibroblast cell
	migration. However, in the case of leukocyte chemotaxis, a smaller pore size (3 $\mu m$ ) is
	recommended.
Components:	1. 96-well Cell Migration Plate : One sterile 96-well plate (see Figure 1 for components)
	2. 96-well Cell Invasion Plate: One sterile 96-well plate containing ECM-coated inserts (see

Figure 1 for components)

- 3. 96-well Cell Harvesting Tray: Two 96-well trays
- 4. Cell Detachment Solution: Two 20 mL bottles
- 5. 4X Lysis Buffer: Two 10 mL bottles
- 6. CyQuant® GR Dye: Two 75  $\mu$ L tubes Top Plate Cover Middle Migration Plate Membrane Chamber Bottom Feeder Tray Figure 1: Components of the 96-well Cell Migration Plate.

#### Material not included:

- 1. Migratory cell lines
- 2. Cell culture medium 3
- 3. Serum free medium, such as DMEM containing 0.5 % BSA, 2 mM CaCl2 and 2 mM MgCl2
- 4. FBS or desired chemoattractant
- 5. Cell culture incubator (37 °C, 5 % CO2 atmosphere)
- 6. Light microscope
- 7. 96-well plate suitable for a fluorescence plate reader
- 8. Fluorescence plate reader

#### **Target Details**

#### Background:

Cell migration is a highly integrated, multistep process that orchestrates embryonic morphogenesis, tissue repair and regeneration. It plays a pivotal role in the disease progression of cancer, mental retardation, atherosclerosis, and arthritis. The initial response of a cell to a migration-promoting agent is to polarize and extend protrusions in the direction of the attractant, these protrusions can consist of large, broad lamellipodia or spike-like filopodia. In either case, these protrusions are driven by actin polymerization and can be stabilized by extracellular matrix (ECM) adhesion or cell-cell interactions. The ability of malignant tumor cells to invade normal surrounding tissue contributes in large part to the significant morbidity and mortality of cancers. Invasiveness requires several distinct cellular functions including adhesion, motility, detachment, and extracellular matrix proteolysis. Metastatic cells produce many proteolytic enzymes (e.g. lysosomal hydrolysates, collagenases, plasminogen activators) while the expression of certain cell surface protease receptors is also increased.

#### **Application Details**

Application Notes:	Optimal working dilution should be determined by the investigator.
Comment:	<ul> <li>Fully quantify chemotaxis and cell invasion with no manual cell counting</li> <li>Includes two plates with 8 µm membrane inserts: one uncoated for chemotaxis and one precoated on top of the membrane with ECM matrix (basement membrane) for cell invasion</li> </ul>
Plate:	Pre-coated
Restrictions:	For Research Use only

#### Handling

Storage:	4 °C
Storage Comment:	Store all components at 4°C.

#### **Publications**

Product cited in:

Pengjam, Madhyastha, Madhyastha, Yamaguchi, Nakajima, Maruyama: "Anthraquinone Glycoside Aloin Induces Osteogenic Initiation of MC3T3-E1 Cells: Involvement of MAPK Mediated Wnt and Bmp Signaling." in: **Biomolecules & therapeutics**, Vol. 24, Issue 2, pp. 123-31 , (2016) (PubMed).

Choi, Lee, Yang, Oh, Won, Han, Jeong, Kim, Kim, Kim, Cho: "Efficient mRNA delivery with graphene oxide-polyethylenimine for generation of footprint-free human induced pluripotent stem cells." in: **Journal of controlled release : official journal of the Controlled Release Society**, Vol. 235, pp. 222-35, (2016) (PubMed).

Lee, Chae, Park, Kim: "Porcine placenta hydrolysates enhance osteoblast differentiation through their antioxidant activity and effects on ER stress." in: **BMC complementary and alternative medicine**, Vol. 16, Issue 1, pp. 291, (2016) (PubMed).

Jin, St Hilaire, Huang, Yang, Dmitrieva, Negro, Schwartzbeck, Liu, Yu, Walts, Davaine, Lee, Donahue, Hsu, Chen, Cheng, Gahl, Chen, Boehm: "Increased activity of TNAP compensates for reduced adenosine production and promotes ectopic calcification in the genetic disease ACDC." in: **Science signaling**, Vol. 9, Issue 458, pp. ra121, (2016) (PubMed).

Pan, Yue, Zhang, Hakim, Kodippili, McDonald, Duan: "AAV-8 is more efficient than AAV-9 in transducing neonatal dog heart." in: **Human gene therapy methods**, Vol. 26, Issue 2, pp. 54-61, (2015) (PubMed).

There are more publications referencing this product on: Product page