

Datasheet for ABIN2345118

HDL Purification Kit (Ultracentrifugation Free)



[Go to Product page](#)

1 Publication

Overview

Quantity:	10 preparations
Application:	Purification (Purif)

Product Details

Purpose:	The HDL Purification Kit uses Dextran Sulfate to selectively precipitate HDL.
Sample Type:	Plasma, Serum
Characteristics:	The kit allows for the purification of HDL without the need for ultracentrifugation. The lipoprotein particles are highly purified through a series of precipitation and low speed centrifugation steps. Each kit provides sufficient reagents to perform up to 10 preps, and each preparation can purify up to 10 mL of serum or plasma samples with a yield of ~4500 µg of HDL per mL for human samples (expected yield will vary by species).
Components:	<ol style="list-style-type: none">1. Dextran Solution : One 8 mL bottle2. Precipitation Solution A : One 30 mL amber bottle3. Tris Solution : One 50 mL bottle containing 20 mM Tris, pH 7.54. Dextran Removal Solution : One 10 mL bottle5. 5X HDL Wash Solution : One 12 mL bottle
Material not included:	<ol style="list-style-type: none">1. Serum or Plasma Samples2. PBS3. Microcentrifuge or Centrifuge4. 10 µL to 1000 µL adjustable single channel micropipettes with disposable tips

Target Details

Background:	Lipoproteins are submicroscopic particles composed of lipid and protein held together by
-------------	--

noncovalent forces. Their general structure is that of a putative spheroidal microemulsion formed from an outer layer of phospholipids, unesterified cholesterol, and proteins, with a core of neutral lipids, predominately cholesteryl esters and triacylglycerols (TAG). Very low density lipoprotein (VLDL), a spherical particle with a diameter of 30-100 nm, is the major plasma vehicle for TAG and is the precursor to Low density lipoprotein (LDL). Each VLDL contains one molecule of a hydrophobic protein known as apolipoprotein B-100 (Apo B), as well as multiple copies of apolipoprotein E and apolipoprotein C. High Density Lipoprotein (HDL) is also a spherical particle with diameter of about 10 nm (Figure 1). HDL contains the Apolipoprotein AI and All molecules. HDL and LDL cholesterol levels in the blood are important indicators of many disease states. High blood levels of LDLs are associated with health problems and cardiovascular disease. For this reason, LDL is often referred to as the "bad cholesterol." LDL particles that accumulate within arteries can form plaques over time, which can increase chances of a stroke, heart attack, or vascular disease. HDL particles are able to remove cholesterol from within arteries and transport it back to the liver for re-utilization or excretion, which is the main reason why the cholesterol carried within HDL particles is sometimes called "good cholesterol." Monitoring circulatory levels of different lipoproteins is critical to the diagnosis of lipid transport disorders such as atherosclerosis. Figure 1: Structure of HDL.

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Reagent Preparation:	<ul style="list-style-type: none">• HDL Resuspension Buffer: Dilute Dextran Solution 1:100 and Precipitation Solution A 1:10 in Tris Solution. For example, add 50 µL of Dextran Solution and 0.5 mL of Precipitation Solution A to 4.45 mL of Tris Solution. Stir to homogeneity. Prepare only enough for immediate use and do not store unused buffer. 3• 1X HDL Wash Solution: Dilute the 5X HDL Wash Solution to 1X with deionized water. Stir to homogeneity. Store unused solution at 4 °C. Purification Protocol Note: The purification protocol below is written for a 10 mL sample size. For smaller sample volumes, scale down each step proportionally. I. Dextran Precipitation 1. To 10 mL of serum or plasma on ice, add 50 µL of Dextran Solution and 500 µL of Precipitation Solution A. Incubate 5 minutes on ice. 2. Spin at 6000 x g 10 minutes at 4 °C. 3. Transfer the supernatant to a new tube for use in section II below. Discard the pellet. II. HDL Purification 1. To 10 mL of supernatant from section I above, add 600 µL of Dextran Solution and 1.5 mL of Precipitation Solution A. Incubate for 2 hours at room temperature. 2. Spin 18,000-20,000 x g for 30 minutes at 4 °C. 3. Discard supernatant and resuspend pellet in 5 mL of HDL Resuspension Buffer (see Preparation of Reagents Section). Mix thoroughly by pipetting up and down. 4. Spin 6000 x g for 10 minutes at 4 °C. 5. Discard supernatant and resuspend pellet in 6 mL of 1X HDL Wash Solution (see Preparation of Reagents Section). 6. Shake for 30 minutes at 4 °C. Shaking speed should be sufficient to dissolve pellet, but not so vigorous that bubbles form. 7. Spin

Application Details

6000 x g for 10 minutes at 4 °C. 8. Transfer the supernatant to a new tube and add 900 µL of Dextran Removal Solution. Mix thoroughly by pipetting up and down. 9. Incubate for 1 hour at 4 °C. 10. Spin 6000 x g for 10 minutes at 4 °C. 11. Transfer the supernatant (containing purified HDL) to a new tube. 12. Dialyze the purified HDL in PBS and determine the protein concentration. 4

Restrictions: For Research Use only

Handling

Storage: 4 °C

Storage Comment: Upon receipt store Dextran Removal Solution at room temperature. Store all other components at 4°C.

Publications

Product cited in: Holzer, Kern, Birner-Grünberger, Curcic, Heinemann, Marsche: "Refined purification strategy for reliable proteomic profiling of HDL2/3: Impact on proteomic complexity." in: **Scientific reports**, Vol. 6, pp. 38533, (2016) ([PubMed](#)).