.-online.com antibodies

Datasheet for ABIN238049 anti-Relaxin 2 antibody



Overview

| Quantity: | 10 µg |
|-----------------------------|--|
| Target: | Relaxin 2 (RLN2) |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Relaxin 2 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Radioimmunoassay (RIA) |
| Product Details | |
| Immunogen: | Purified human Relaxin-2 |
| Cross-Reactivity (Details): | Does not cross-react with Insulin, Zn-Insulin, IGF-1, IGF-2, Spermolaxin, Inhibin alpha subunit, |
| | Inhibin, Seminal-Plasma-Inhibin-like peptide, CG, LH, FSH, Prolactin and Ubiquitin. |
| Characteristics: | Rabbit anti Human Relaxin-2, Rabbit antibody to Human Relaxin-2 |
| Purification: | Protein G |
| | |
| | |

| Target: | Relaxin 2 (RLN2) |
|-------------------|---------------------------|
| Alternative Name: | Relaxin-2 (RLN2 Products) |
| Pathways: | Hormone Activity |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN238049 | 09/12/2023 | Copyright antibodies-online. All rights reserved.

| Application Details | |
|---------------------|---|
| Application Notes: | Suitable for use in ELISA (1 μ g/mL), RIA (1 μ g/mL) and Immunohistochemistry (5 μ g/mL). Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Lyophilized |
| Reconstitution: | Reconstitute in 10 µL double distilled water. |
| Buffer: | Lyophilized from 50 mM Tris, pH 7.5. |
| Preservative: | Without preservative |
| Handling Advice: | Avoid multiple freeze/thaw cycles. |
| Storage: | 4 °C/-20 °C |
| Storage Comment: | Store lyophilized product at 2-8 °C. After reconstitution, store at -28 °C. |