antibodies -online.com





anti-MAL antibody (Internal Region)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μg
Target:	MAL
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This MAL antibody is un-conjugated
Application:	ELISA

Product Details

Purpose:	T-cell differentiation protein	
Immunogen:	Peptide with sequence C-QDGFTYRHYHEN, from the internal region of the protein sequence according to NP_002362.1, NP_071884.1.	
Sequence:	QDGFTYRHYH EN	
Isotype:	IgG	
Specificity:	This antibody is expected to recognise isoform a (NP_002362.1) and isoform c (NP_071884.1).	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.	
Grade:	Recent	

Target Details

rarget Details		
Target:	MAL	
Alternative Name:	T-cell differentiation protein (MAL Products)	
Background:	MAL, mal, T-cell differentiation protein , T-cell differentiation protein MAL, T-lymphocyte	
	maturation-associated protein, myelin and lymphocyte protein	
Gene ID:	4118, 17153, 25263	
NCBI Accession:	NP_002362, NP_071884	
Pathways:	Asymmetric Protein Localization	
Application Details		
Application Notes:	Western Blot: Preliminary experiments gave an approx 35 kDa band in Human Thymus and	
	Thyroid Gland lysates after 0.3 μ g/mL antibody staining. Please note that currently we cannot	
	find an explanation in the literature for the band we observe given the calcu	
	Peptide ELISA: antibody detection limit dilution 1:4000.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	0.5 mg/mL	
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum	
	albumin.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Handling Advice:	Minimize freezing and thawing.	
Storage:	-20 °C	
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerate	
	at 4°C for a few weeks and still remain viable.	