
antibodies - online.com







anti-Actl7b antibody (Internal Region)



Image



Overview

Quantity:	100 μg
Target:	Actl7b
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This ActI7b antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Purpose:	Actin-like 7B
Immunogen:	Peptide with sequence C-QLKMKPRKVHKIK, from the internal region of the protein sequence according to NP_006677.1.
Sequence:	QLKMKPRKVH KIK
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

Target Details

rarget Details	
Target:	Actl7b
Alternative Name:	ACTL7B (Actl7b Products)
Background:	Actin-like 7B, ACTL7B, RP11-3J11.2, actin-like 7-beta
Gene ID:	10880
NCBI Accession:	NP_006677
Application Details	
Application Notes:	Western Blot: Approx 48 kDa band observed in human testis lysates (calculated MW of 45.2 kDa according to NP_006677.1). Recommended concentration: 0.03-0.1 µg/mL.
	Peptide ELISA: antibody detection limit dilution 1:128000.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

15kDa

Image 1. ABIN238621 ($0.03\mu g/ml$) staining of human testis lysate ($35\mu g$ protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.