



[Go to Product page](#)

Datasheet for ABIN2452175

SSB Protein

3 Images

2 Publications

Overview

Quantity:	200 µg
Target:	SSB
Origin:	E. coli
Source:	Escherichia coli (E. coli)
Protein Type:	Native
Biological Activity:	Active
Application:	Functional Studies (Func)

Product Details

Purity: > 95 % purity as determined by SDS-PAGE (CBB staining)

Target Details

Target:	SSB
Abstract:	SSB Products
Background:	E.coli single-stranded DNA binding protein (SSB) binds to single-stranded DNA with high specificity. It is involved in DNA replication and recombination in vivo. The SSB gene was expressed as the recombinant protein in E.coli highly purified. The molecular mass is 18.9 kDa.
UniProt:	P0AGE0

Application Details

Application Notes: 1. Functional single-stranded DNA-binding protein for studying DNA replication and

Application Details

recombination

2. Enhancement of the specificity and yield of PCR

Comment: The absence of endonucleases and exonucleases was confirmed.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 5 mg/mL

Buffer: 20 mM Tris-HCl (pH 7.6), 200 mM NaCl, 1 mM dithiothreitol, 1 mM EDTA, 50 % glycerol

Preservative: Dithiothreitol (DTT)

Precaution of Use: This product contains Dithiothreitol (DTT): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Publications

Product cited in: Aaronson, Bottaro, Miki, Ron, Finch, Fleming, Ahn, Taylor, Rubin: "Keratinocyte growth factor. A fibroblast growth factor family member with unusual target cell specificity." in: **Annals of the New York Academy of Sciences**, Vol. 638, pp. 62-77, (1992) ([PubMed](#)).

Rubin, Osada, Finch, Taylor, Rudikoff, Aaronson: "Purification and characterization of a newly identified growth factor specific for epithelial cells." in: **Proceedings of the National Academy of Sciences of the United States of America**, Vol. 86, Issue 3, pp. 802-6, (1989) ([PubMed](#)).





Fig.1 SDS-PAGE of *E.coli* SSB protein

Western Blotting


Image 1.



Polymerase Chain Reaction

Image 2.

0 1 2 3



0.02 ug/ul of M13mp18ssDNA was incubated with 0 (lane 0), 0.025 (lane 1), 0.05 (lane 2), and 0.1(lane 3) ug/ul of SSB at 37 °C for 30 min. and then 10ul aliquot was subjected to electrophoresis in agarose.

Polymerase Chain Reaction

Image 3.