antibodies -online.com







Rat Normal Tissue Blot II



Image



Overview

Quantity:	1 applications
Host:	Rat
Application:	Western Blotting (WB)

Product Details	
Characteristics:	The rat tissue blot is designed for proteomics research in screening for antibody and protein
	expression. It carries 8 different rat tissue lysates with 15 µg of total cellular protein in each
	lane. Proteins were separated in 4-20 % SDS-PAGE gel and transferred onto nitrocellulose
	membrane. The rat tissues included in the blot are adrenal, cerebellum, cerebrum, small
	intestine, skeletal muscle, stomach, testis, and thymus.
	Sample diagnosis: Normal
	Samples loaded: Lanes: 1: protein markers, 2: rat rat adrenal, 3: cerebellum, 4: cerebrum, 5:
	small intestine, 6: skeletal muscle, 7: stomach, 8: testis, and 9: thymus
	Quantity loaded per lane: 15 μg

Application Details


Application Details	
Application Notes:	For Western Blot, the blot can be blocked with 1 % BSA or 5 % nonfat dry milk for 1 hour.
	Incubation takes 30 minutes to two hours with primary antibody, and 30 minutes to one hour
	with secondary antibody at room temperature.
Restrictions:	For Research Use only

Gel percentage: Gradient 4-20 %

Handling

Handling Advice:	The blot can be reused after stripping.
Storage:	4 °C
Storage Comment:	Store at 4 °C.

Images



SDS-PAGE

Image 1.