antibodies -online.com





anti-OR6C75 antibody



Image



Go to Product page

\sim							
	1//	$rac{1}{2}$	`\/	L	Р	1	٨

100 μL	
OR6C75	
Human, Rat, Mouse	
Rabbit	
Polyclonal	
This OR6C75 antibody is un-conjugated	
Western Blotting (WB), ELISA	
Antibody produced in rabbits immunized with a synthetic peptide corresponding a region of human OR6C75.	
Antibody is purified by peptide affinity chromatography method.	
OR6C75	
OR6C75 (OR6C75 Products)	

transduction of odorant signals. The olfactory receptor gene family is the largest in the
genome. The nomenclature assigned to the olfactory receptor genes and proteins for this
organism is independent of other organisms.Olfactory receptors interact with odorant
molecules in the nose, to initiate a neuronal response that triggers the perception of a smell.
The olfactory receptor proteins are members of a large family of G-protein-coupled receptors
(GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane
domain structure with many neurotransmitter and hormone receptors and are responsible for
the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor
gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor
genes and proteins for this organism is independent of other organisms.
PRIMARYREFSEQ_SPAN PRIMARY_IDENTIFIER PRIMARY_SPAN COMP 1-939 AC125816.8
75932-76870 c

Molecular Weight:	35 kDa
Gene ID:	390323
NCBI Accession:	NP_001005497
UniProt:	A6NL08

Application Details

Application Notes:	OR6C75 antibody can be used for detection of OR6C75 by ELISA at 1:312500. OR6C75 antibody

can be used for detection of OR6C75 by western blot at 1 µg/mL, and HRP conjugated

secondary antibody should be diluted 1:50,000 - 100,000.


Restrictions: For Research Use only

Handling

Format:	Lyophilized	
Reconstitution:	Add 50 ?L of distilled water. Final antibody concentration is 1 mg/mL.	
Concentration:	ation: 1 mg/mL	
Buffer:	Antibody is lyophilized in PBS buffer with 2 % sucrose.	
Handling Advice:	As with any antibody avoid repeat freeze-thaw cycles.	
Storage:	4 °C/-20 °C	
Storage Comment:	For short periods of storage (days) store at 4 °C. For longer periods of storage, store OR6C75	

antibody at -20 °C.

Images



Western Blotting

Image 1.