antibodies -online.com





anti-CAND2 antibody



Image



Go to Product page

| \sim | | | |
|--------|-----|------|------------|
| | N/6 | 1//r | $I \cap V$ |

| Quantity: | 100 μL |
|-------------------|---|
| Target: | CAND2 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This CAND2 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA |
| Product Details | |
| Immunogen: | Antibody produced in rabbits immunized with a synthetic peptide corresponding a region of |
| | human CAND2. |
| Purification: | Antibody is purified by peptide affinity chromatography method. |
| Target Details | |
| Target: | CAND2 |
| Alternative Name: | CAND2 (CAND2 Products) |
| Background: | CAND2 may play a role in the assembly of ubiquitin ligase complexes and modulate the |
| | ubiquitination of target proteins. And it may be also a transcription regulator. |
| Molecular Weight: | 123 kDa |
| Gene ID: | 23066 |
| | |

Target Details

| NCBI Accession: | NP_036430 |
|-----------------|-----------|
| UniProt: | 075155 |

Application Details


| Application Notes: | CAND2 antibody can be used for detection of CAND2 by ELISA at 1:62500. CAND2 antibody can | |
|--------------------|--|--|
| | be used for detection of CAND2 by western blot at 1 μ g/mL, and HRP conjugated secondary | |
| | antibody should be diluted 1:50,000 - 100,000. | |

Restrictions: For Research Use only

Handling

| Format: | Lyophilized | |
|------------------|---|--|
| Reconstitution: | Add 50 ?L of distilled water. Final antibody concentration is 1 mg/mL. | |
| Concentration: | 1 mg/mL | |
| Buffer: | Antibody is lyophilized in PBS buffer with 2 % sucrose. | |
| Handling Advice: | As with any antibody avoid repeat freeze-thaw cycles. | |
| Storage: | 4 °C/-20 °C | |
| Storage Comment: | For short periods of storage (days) store at 4 °C. For longer periods of storage, store CAND2 | |
| | antibody at -20 °C. | |

Images



Western Blotting

Image 1.