
antibodies -online.com





anti-SNW1 antibody





Overview	
Quantity:	100 μL
Target:	SNW1
Reactivity:	Human, Dog, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SNW1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA
Product Details	
Immunogen:	Antibody produced in rabbits immunized with a synthetic peptide corresponding a region of
	human SKIIP.
Purification:	Antibody is purified by peptide affinity chromatography method.
Target Details	
Target:	SNW1
Alternative Name:	SKIIP (SNW1 Products)
Background:	SKIIP (Nuclear Protein SkiP, nuclear receptor coactivator NCoA-62, ski-interacting protein) is a
	member of the SNW gene family, encodes a coactivator that enhances transcription from some
	Pol II promoters. This coactivator can bind to the ligand-binding domain of the vitamin D
	receptor and to retinoid receptors to enhance vitamin D-, retinoic acid-, estrogen-, and

glucocorticoid-mediated gene expression. It can also interact with poly (A)-binding protein 2 to


Target Details

Storage Comment:

	directly control the expression of muscle-specific genes at the transcriptional level. Finally, the protein may be involved in oncogenesis since it interacts with a region of SKI oncoproteins that is required for transforming activity.
Molecular Weight:	61 kDa
Gene ID:	22938
NCBI Accession:	NP_036377
UniProt:	Q13573
Pathways:	Retinoic Acid Receptor Signaling Pathway
Application Details	
Application Notes:	SKIIP antibody can be used for detection of SKIIP by ELISA at 1:62500. SKIIP antibody can be used for detection of SKIIP by western blot at 1.0-5.0 μ g/mL, and HRP conjugated secondary antibody should be diluted 1:50,000 - 100,000.
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Add 50 ?L of distilled water. Final antibody concentration is 1 mg/mL.
Concentration:	1 mg/mL
Buffer:	Antibody is lyophilized in PBS buffer with 2 % sucrose.
Handling Advice:	As with any antibody avoid repeat freeze-thaw cycles.
Storage:	4 °C/-20 °C

antibody at -20 °C.

For short periods of storage (days) store at 4 °C. For longer periods of storage, store SKIIP



Western Blotting

Image 1.