antibodies -online.com





Datasheet for ABIN2461507

anti-TSC22D4 antibody





| \sim | | |
|-----------------------|-------|-------------------|
| $\bigcap \bigvee \in$ | K\ /I | $\triangle V V V$ |
| | | |

| Overview | |
|-----------------|-------------------------------------------------------------------------------------------|
| Quantity: | 100 μL |
| Target: | TSC22D4 |
| Reactivity: | Rat, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This TSC22D4 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (IHC) |
| Product Details | |
| Immunogen: | Antibody produced in rabbits immunized with a synthetic peptide corresponding a region of |

| Immunogen: | Antibody produced in rabbits immunized with a synthetic peptide corresponding a region of | |
|---------------|-------------------------------------------------------------------------------------------|--|
| | mouse Tsc22d4. | |
| Purification: | Antibody is purified by protein A chromatography method. | |

Target Details

| Target: | TSC22D4 |
|-------------------|---------------------------------------------|
| Alternative Name: | Tsc22d4 (TSC22D4 Products) |
| Background: | Tsc22d4 may be a transcriptional repressor. |
| Molecular Weight: | 43 kDa |
| Gene ID: | 78829 |
| NCBI Accession: | NP_076399 |

Target Details

UniProt: Q9EQN3

Application Details

Application Notes: Tsc22d4 antibody can be used for detection of Tsc22d4 by ELISA at 1:62500. Tsc22d4

antibody can be used for detection of Tsc22d4 by western blot at 5.0 µg/mL, and HRP


conjugated secondary antibody should be diluted 1:50,000 - 100,000.

Restrictions: For Research Use only

Handling

| Format: | Lyophilized |
|------------------|-------------------------------------------------------------------------------------------------|
| Reconstitution: | Add 50 ?L of distilled water. Final antibody concentration is 1 mg/mL. |
| Concentration: | 1 mg/mL |
| Buffer: | Antibody is lyophilized in PBS buffer with 2 % sucrose. |
| Handling Advice: | As with any antibody avoid repeat freeze-thaw cycles. |
| Storage: | 4 °C/-20 °C |
| Storage Comment: | For short periods of storage (days) store at 4 °C. For longer periods of storage, store Tsc22d4 |
| | antibody at -20 °C. |

Images



Western Blotting

Image 1.