antibodies -online.com





anti-ANAPC7 antibody





| _ | | | | | | |
|---|---|----|---------|---|---|---|
| 0 | V | е. | r_{V} | / | 6 | W |

| Quantity: | 100 μL |
|-------------------|---|
| Target: | ANAPC7 |
| Reactivity: | Human, Mouse, Dog |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This ANAPC7 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (IHC) |
| Product Details | |
| Immunogen: | Antibody produced in rabbits immunized with a synthetic peptide corresponding a region of |
| | human ANAPC7. |
| Purification: | Antibody is purified by peptide affinity chromatography method. |
| Target Details | |
| Target: | ANAPC7 |
| Alternative Name: | ANAPC7 (ANAPC7 Products) |
| Background: | The anaphase-promoting complex (APC) consists of at least 8 protein subunits, including |
| | APC5, CDC27 (APC3), CDC16 (APC6), and CDC23 (APC8). The anaphase-promoting complex |
| | (APC) consists of at least 8 protein subunits, including APC5, CDC27 (APC3, MIM 116946), |
| | CDC16 (APC6, MIM 603461), and CDC23 (APC8, MIM 603462). |
| Molecular Weight: | 62 kDa |
| | |

Target Details

| Gene ID: | 51434 |
|-----------------|-----------|
| NCBI Accession: | NP_057322 |
| UniProt: | Q9UJX3 |


Application Details

| Application Notes: | ANAPC7 antibody can be used for detection of ANAPC7 by ELISA at 1:62500. ANAPC7 antibody |
|--------------------|--|
| | can be used for detection of ANAPC7 by western blot at 0.5 $\mu g/mL$, and HRP conjugated |
| | secondary antibody should be diluted 1:50,000 - 100,000. |
| Restrictions: | For Research Use only |

Handling

| Format: | Lyophilized | |
|------------------|--|--|
| Reconstitution: | Add 50 ?L of distilled water. Final antibody concentration is 1 mg/mL. | |
| Concentration: | 1 mg/mL | |
| Buffer: | Antibody is lyophilized in PBS buffer with 2 % sucrose. | |
| Handling Advice: | As with any antibody avoid repeat freeze-thaw cycles. | |
| Storage: | 4 °C/-20 °C | |
| Storage Comment: | For short periods of storage (days) store at 4 °C. For longer periods of storage, store ANAPC7 | |
| | antibody at -20 °C. | |

Images



Western Blotting

Image 1.