antibodies -online.com





anti-TIGD3 antibody






Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL	
Target:	TIGD3	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TIGD3 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA	
Product Details		
Immunogen:	Antibody produced in rabbits immunized with a synthetic peptide corresponding a region of	
	human TIGD3.	
Purification:	Antibody is purified by protein A chromatography method.	
Target Details		
Target:	TIGD3	
Alternative Name:	TIGD3 (TIGD3 Products)	
Background:	TIGD3 belongs to the tigger subfamily of the pogo superfamily of DNA-mediated transposons in	
	humans. These proteins are related to DNA transposons found in fungi and nematodes, and	
	more distantly to the Tc1 and mariner transposases. They are also very similar to the major	
	mammalian centromere protein B. The exact function of TIGD3 gene is not known. The protein	
	encoded by this gene belongs to the tigger subfamily of the pogo superfamily of DNA-mediated	

Target Details		
	transposons in humans. These proteins are related to DNA transposons found in fungi and nematodes, and more distantly to the Tc1 and mariner transposases. They are also very similar to the major mammalian centromere protein B. The exact function of this gene is not known.	
Molecular Weight:	52 kDa	
Gene ID:	220359	
NCBI Accession:	NP_663771	
UniProt:	Q6B0B8	
Application Details		
Application Notes:	TIGD3 antibody can be used for detection of TIGD3 by ELISA at 1:1562500. TIGD3 antibody can be used for detection of TIGD3 by western blot at 1.25 µg/mL, and HRP conjugated secondary antibody should be diluted 1:50,000 - 100,000.	
Restrictions:	For Research Use only	
Handling		
Format:	Lyophilized	
Reconstitution:	Add 100 ?L of distilled water. Final antibody concentration is 1 mg/mL.	
Concentration:	1 mg/ml	

Format:	Lyophilized	
Reconstitution:	Add 100 ?L of distilled water. Final antibody concentration is 1 mg/mL.	
Concentration:	1 mg/mL	
Buffer:	Antibody is lyophilized in PBS buffer with 2 % sucrose.	
Handling Advice:	As with any antibody avoid repeat freeze-thaw cycles.	
Storage:	4 °C/-20 °C	
Storage Comment:	For short periods of storage (days) store at 4 °C. For longer periods of storage, store TIGD3 antibody at -20 °C.	



Western Blotting

Image 1.