antibodies .- online.com







anti-Very Low Density Lipoprotein (VLDL) antibody



Image



\sim	
()\/\	rview
	VICVV

Restrictions:


effect of glucocorticoids on arachidonic acid release and prostaglandin I2 formation in va smooth muscle cells. Application Details	OVERVIEW	
Reactivity: Human Host: Chicken Clonality: Polyclonal Application: Western Blotting (WB), ELISA Product Details Purification: Immunoaffinity Purified Target Details Target: Very Low Density Lipoprotein (VLDL) Alternative Name: VLDL (VLDL Products) Background: VLDL transports hepatic synthesized triglycerides and cholesterol. May counteract inhibite effect of glucocorticoids on arachidonic acid release and prostaglandin I2 formation in vasmooth muscle cells. Application Details Application Notes: VLDL antibody can be used for the detection of VLDL by ELISA and Western Blot, may also	Quantity:	0.05 mg
Clonality: Polyclonal Application: Western Blotting (WB), ELISA Product Details Purification: Immunoaffinity Purified Target Details Target: Very Low Density Lipoprotein (VLDL) Alternative Name: VLDL (VLDL Products) Background: VLDL transports hepatic synthesized triglycerides and cholesterol. May counteract inhibit effect of glucocorticoids on arachidonic acid release and prostaglandin 12 formation in vasmooth muscle cells. Application Details Application Notes: VLDL antibody can be used for the detection of VLDL by ELISA and Western Blot, may also	Target:	Very Low Density Lipoprotein (VLDL)
Clonality: Polyclonal Application: Western Blotting (WB), ELISA Product Details Purification: Immunoaffinity Purified Target Details Target: Very Low Density Lipoprotein (VLDL) Alternative Name: VLDL (VLDL Products) Background: VLDL transports hepatic synthesized triglycerides and cholesterol. May counteract inhibite effect of glucocorticoids on arachidonic acid release and prostaglandin 12 formation in vasmooth muscle cells. Application Details Application Notes: VLDL antibody can be used for the detection of VLDL by ELISA and Western Blot, may also	Reactivity:	Human
Application: Western Blotting (WB), ELISA Product Details Purification: Immunoaffinity Purified Target Details Target: Very Low Density Lipoprotein (VLDL) Alternative Name: VLDL (VLDL Products) Background: VLDL transports hepatic synthesized triglycerides and cholesterol. May counteract inhibit effect of glucocorticoids on arachidonic acid release and prostaglandin 12 formation in vasmooth muscle cells. Application Details Application Notes: VLDL antibody can be used for the detection of VLDL by ELISA and Western Blot, may also	Host:	Chicken
Purification: Immunoaffinity Purified Target Details Target: Very Low Density Lipoprotein (VLDL) Alternative Name: VLDL (VLDL Products) Background: VLDL transports hepatic synthesized triglycerides and cholesterol. May counteract inhibit effect of glucocorticoids on arachidonic acid release and prostaglandin 12 formation in vasmooth muscle cells. Application Details Application Notes: VLDL antibody can be used for the detection of VLDL by ELISA and Western Blot, may also	Clonality:	Polyclonal
Purification: Immunoaffinity Purified Target Details Target: Very Low Density Lipoprotein (VLDL) Alternative Name: VLDL (VLDL Products) Background: VLDL transports hepatic synthesized triglycerides and cholesterol. May counteract inhibite effect of glucocorticoids on arachidonic acid release and prostaglandin I2 formation in vasmooth muscle cells. Application Details Application Notes: VLDL antibody can be used for the detection of VLDL by ELISA and Western Blot, may also	Application:	Western Blotting (WB), ELISA
Target: Very Low Density Lipoprotein (VLDL) Alternative Name: VLDL (VLDL Products) Background: VLDL transports hepatic synthesized triglycerides and cholesterol. May counteract inhibit effect of glucocorticoids on arachidonic acid release and prostaglandin I2 formation in value smooth muscle cells. Application Details Application Notes: VLDL antibody can be used for the detection of VLDL by ELISA and Western Blot, may also	Product Details	
Target: Very Low Density Lipoprotein (VLDL) Alternative Name: VLDL (VLDL Products) Background: VLDL transports hepatic synthesized triglycerides and cholesterol. May counteract inhibit effect of glucocorticoids on arachidonic acid release and prostaglandin I2 formation in vasmooth muscle cells. Application Details Application Notes: VLDL antibody can be used for the detection of VLDL by ELISA and Western Blot, may also	Purification:	Immunoaffinity Purified
Alternative Name: VLDL (VLDL Products) Background: VLDL transports hepatic synthesized triglycerides and cholesterol. May counteract inhibit effect of glucocorticoids on arachidonic acid release and prostaglandin I2 formation in vasmooth muscle cells. Application Details Application Notes: VLDL antibody can be used for the detection of VLDL by ELISA and Western Blot, may also	Target Details	
Background: VLDL transports hepatic synthesized triglycerides and cholesterol. May counteract inhibite effect of glucocorticoids on arachidonic acid release and prostaglandin I2 formation in value smooth muscle cells. Application Details VLDL antibody can be used for the detection of VLDL by ELISA and Western Blot, may also	Target:	Very Low Density Lipoprotein (VLDL)
effect of glucocorticoids on arachidonic acid release and prostaglandin I2 formation in value smooth muscle cells. Application Details Application Notes: VLDL antibody can be used for the detection of VLDL by ELISA and Western Blot, may also	Alternative Name:	VLDL (VLDL Products)
Application Notes: VLDL antibody can be used for the detection of VLDL by ELISA and Western Blot, may also	Background:	VLDL transports hepatic synthesized triglycerides and cholesterol. May counteract inhibitory effect of glucocorticoids on arachidonic acid release and prostaglandin I2 formation in vascular smooth muscle cells.
	Application Details	
	Application Notes:	VLDL antibody can be used for the detection of VLDL by ELISA and Western Blot, may also work for ICC and IHC.

For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Phosphate-Buffered Saline. No preservatives added.
Preservative:	Without preservative
Handling Advice:	As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Storage:	4 °C/-20 °C
Storage Comment:	VLDL antibody can be stored at 4 °C for short term (weeks). Long term storage should be at -20 °C.

Images



ELISA

Image 1.