

Datasheet for ABIN2469321

anti-DDR1 antibody (C-Term)





Go to Product page

| 0 | | | | |
|---|--|--|--|--|
| | | | | |
| | | | | |
| | | | | |


| Quantity: | 0.05 mg | | |
|---|---|--|--|
| Target: | DDR1 | | |
| Binding Specificity: | C-Term | | |
| Reactivity: | Human, Monkey, Pig, Bat, Horse, Cow, Dog, Chimpanzee, Gibbon, Gorilla | | |
| Host: | Rabbit | | |
| Clonality: | Polyclonal | | |
| Conjugate: | This DDR1 antibody is un-conjugated | | |
| Application: | Immunohistochemistry (IHC) | | |
| | | | |
| Product Details | | | |
| Product Details Immunogen: | DDR1 antibody was raised against a peptide located near the C-Terminal region of DDR1 (Human). | | |
| | | | |
| Immunogen: | (Human). BLAST analysis of the peptide immunogen showed no homology with other human proteins, | | |
| Immunogen: Specificity: | (Human). BLAST analysis of the peptide immunogen showed no homology with other human proteins, except DDR2 (69â€⁻%). | | |
| Immunogen: Specificity: Purification: | (Human). BLAST analysis of the peptide immunogen showed no homology with other human proteins, except DDR2 (69â€⁻%). | | |
| Immunogen: Specificity: Purification: Target Details | (Human). BLAST analysis of the peptide immunogen showed no homology with other human proteins, except DDR2 (69â€⁻%). Immunoaffinity Chromatography | | |

Target Details

Storage Comment:

| rarget Details | | |
|---------------------|--|--|
| | and a delayed but sustained tyrosine kinase activation. DDR1 may function in cell-to-cell interaction or recognition. At least three mRNA variants, resulting in different protein isoforms of 875, 913 and 919 amino acids, have been reported. DDR1 protein has been shown to be over expressed in human breast, ovarian, esophageal and pediatric brain tumors. | |
| Gene ID: | 780 | |
| UniProt: | Q08345 | |
| Pathways: | RTK Signaling, Smooth Muscle Cell Migration | |
| Application Details | | |
| Application Notes: | DDR1 antibody can be used in ELISA, Western Blot starting at $1 - 2 \mu g/mL$, immunocytochemistry, immunohistochemistry starting at $10 \mu g/mL$, immunoprecipitation starting at $1 - 2 \mu g/mL$, and flow cytometry. | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Buffer: | PBS, 0.1 % sodium azide. | |
| Preservative: | Sodium azide | |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. | |
| Handling Advice: | As with all antibodies avoid freeze/thaw cycles. | |
| Storage: | 4 °C/-80 °C | |

DDR1 antibody should be stored long term (months) at -80 °C and short term (days) at 4 °C.



Immunohistochemistry

Image 1.