

Datasheet for ABIN2474276
Goat anti-Dog IgE Antibody



[Go to Product page](#)

1 Publication

Overview

| | |
|--------------|------------------------------|
| Quantity: | 0.5 mg |
| Target: | IgE |
| Reactivity: | Dog |
| Host: | Goat |
| Clonality: | Polyclonal |
| Application: | ELISA, Western Blotting (WB) |

Product Details

| | |
|------------------|--------------|
| Isotype: | IgG |
| Characteristics: | Purified IgG |
| Purification: | Purified |

Target Details

| | |
|-----------|------------------------------|
| Target: | IgE |
| Abstract: | IgE Products |

Application Details

| | |
|--------------------|--|
| Application Notes: | Optimal working dilution should be determined by the investigator. |
| Restrictions: | For Research Use only |

Handling

Format: Liquid

Concentration: 1.0 mg/mL

Publications

Product cited in: Motallebzadeh, Rehakova, Conlon, Win, Callaghan, Goddard, Bolton, Ruddle, Bradley, Pettigrew: "Blocking lymphotoxin signaling abrogates the development of ectopic lymphoid tissue within cardiac allografts and inhibits effector antibody responses." in: **FASEB journal : official publication of the Federation of American Societies for Experimental Biology**, Vol. 26, Issue 1, pp. 51-62, (2012) ([PubMed](#)).

Silveira, Nunes, Cara, Souza, Corrêa, Teixeira, Negrão-Corrêa: "Infection with *Strongyloides venezuelensis* induces transient airway eosinophilic inflammation, an increase in immunoglobulin E, and hyperresponsiveness in rats." in: **Infection and immunity**, Vol. 70, Issue 11, pp. 6263-72, (2002) ([PubMed](#)).