antibodies .- online.com

# Datasheet for ABIN2481645 anti-Calnexin antibody (C-Term) (PE)

5 Images



Overview

Quantity:	200 µg
Target:	Calnexin (CANX)
Binding Specificity:	C-Term
Reactivity:	Dog
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Calnexin antibody is conjugated to PE
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunoprecipitation (IP), Flow Cytometry (FACS), Immunocytochemistry (ICC)
Product Details	
Immunogen:	Dog Calnexin C-terminal synthetic peptide conjugated to KLH. Identical to human, mouse and rat calnexin sequences over these residues.
Specificity:	Detects the C-terminal domain of Calnexin ~90 kDa. Weak detection in Chicken, Drosophila, and

Immunogen:	Dog Calnexin C-terminal synthetic peptide conjugated to KLH. Identical to human, mouse and rat calnexin sequences over these residues.
Specificity:	Detects the C-terminal domain of Calnexin ~90 kDa. Weak detection in Chicken, Drosophila, and Xenopus tissues.
Cross-Reactivity:	Chicken, Cow, Dog, Drosophila melanogaster, Guinea Pig, Hamster, Human, Monkey, Mouse, Pig, Quail, Rabbit, Rat, Sheep, Xenopus laevis
Purification:	Protein A Purified

## Target Details

Target:

Calnexin (CANX)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN2481645 | 09/11/2023 | Copyright antibodies-online. All rights reserved.


Target Details		
Alternative Name:	Calnexin (CANX Products)	
Background:	Calnexin, an abundant ~90 kDa integral protein of the endoplasmic reticulum, is also referred to as IP90, p88 and p90 (1). It consists of a large 50 kDa N-terminal calcium-binding luminal domain, a single transmembrane helix and a short acidic cytoplasmic tail (2, 3). Unlike its ER counterparts which have a KDEL sequence on their C-terminus to ensure ER retention (4), calnexin has positively charged cytosolic residues that do the same thing (3). Most ER proteins act as molecular chaperones and participate in the proper folding of polypeptides and their assembly into multi-subunit proteins. Calnexin together with calreticulin, plays a key role in glycoprotein folding and its control within the ER, by interacting with folding intermediates via their mono-glycosylated glycans (5, 6). Calnexin has also been shown to associate with the major histocompatibility complex class I heavy chains, partial complexes of the T cell receptor and B cell membrane immunoglobulin (7).	
Gene ID:	403908	
NCBI Accession:	NP_001003232	
UniProt:	P24643	
Pathways:	MAPK Signaling, Thyroid Hormone Synthesis	
Application Details		
Application Notes:	<ul> <li>WB (1:2000)</li> <li>ICC/IF (1:100)</li> <li>IHC (1:100)</li> <li>optimal dilutions for assays should be determined by the user.</li> </ul>	
Comment:	A 1:2000 dilution of ABIN2481645 was sufficient for detection of Calnexin in 10 $\mu$ g of HeLa cell lysate by ECL immunoblot analysis.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	PBS pH 7.2, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated	
Preservative:	Sodium azide	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN2481645 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

```
Handling
```

Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Conjugated antibodies should be stored at 4°C

## Images






### Immunofluorescence (fixed cells)

**Image** 1. Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-Calnexin-CT Polyclonal Antibody . Tissue: HaCaT cells. Species: Human. Fixation: Cold 100% methanol at -20C for 10 minutes. Primary Antibody: Rabbit Anti-Calnexin-CT Polyclonal Antibody at 1:100 for 12 hours at 4°C. Secondary Antibody: FITC Goat Anti-Rabbit at 1:50 for 1-2 hours at RT in dark. Localization: Nuclear staining, cytoplasmic staining.

#### Immunofluorescence (fixed cells)

2. Immunocytochemistry/Immunofluorescence Image analysis using Rabbit Anti-Calnexin-CT Polyclonal Antibody . Tissue: Heat Shocked HeLa Cells. Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Rabbit Anti-Calnexin-CT Polyclonal Antibody at 1:80 for 12 hours at 4°C. Secondary Antibody: R-PE Goat Anti-Rabbit (yellow) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Endoplasmic reticulum membrane. Melanosome. Magnification: 20x. (A) DAPI (blue) nuclear stain. (B) Anti-Calnexin-CT Antibody. (C) Composite. Heat Shocked at 42°C for 1h.



#### Immunofluorescence (fixed cells)

**Image 3.** Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-Calnexin-CT Polyclonal Antibody . Tissue: Heat Shocked HeLa Cells. Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Rabbit Anti-Calnexin-CT Polyclonal Antibody at 1:80 for 12 hours at 4°C. Secondary Antibody: FITC Goat Anti-Rabbit (green) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Endoplasmic reticulum membrane. Melanosome. Magnification: 100x. Heat Shocked at 42°C for 1h.

Please check the product details page for more images. Overall 5 images are available for ABIN2481645.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 4/4 | Product datasheet for ABIN2481645 | 09/11/2023 | Copyright antibodies-online. All rights reserved.