antibodies.com

# Datasheet for ABIN2481837 anti-CRYAA antibody (Atto 594)

2 Images



#### Overview

Quantity:	200 µg
Target:	CRYAA
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CRYAA antibody is conjugated to Atto 594
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

## Product Details

Immunogen:	Native Alpha Crystallin	
Clone:	1H3-B8	
lsotype:	lgG1	
Specificity:	Detects ~20 kDa. Does not cross-react with $\alpha B$ -crystallin, $\beta L$ -crystallin, BH- crystallin, $\gamma$ -	
	crystallin, HSP25, HSP27 or HSP47 proteins.	
Cross-Reactivity:	Cow, Human, Mouse, Rat	
Purification:	Protein G Purified	
Target Details		
Target:	CRYAA	

Alternative Name:	Alpha A Crystallin (CRYAA Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN2481837 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

### Target Details

Background:	The alpha-crystallins are major water-soluble lens structural proteins of the vertebrate eye that
	are related to the small heat shock protein family. The alpha-crystallins possess structural and
	functional similarities with HSP25 and HSP27 (1). Mammalian lens cystallins are divided into
	alpha, beta and gamma families. Alpha and beta families are further divided into acidic and
	basic groups (Alpha-A and Alpha-B respectively). In the lens, alpha-crystallin primarily functions
	to maintain proper refractive index, however it can also function as a molecular chaperone that
	binds to the denatured proteins, keeping them in solution and thereby maintaining the
	translucency of the lens. When cellular stress occurs, alpha-crystallin enters its' phosphorylated
	state and may serve a structural control function and play a role in protein maintenance (2). In
	addition to their interaction with proteins, alpha-crystallins also interact with native molecules
	such as membrane proteins, Golgi matrix protein, structural proteins, nuclear proteins and DNA
	(3, 4, 5, 6, and 7). Two other functions are an autokinase activity and participation in the
	intracellular architecture, and it has also been proven that both alpha-A and B prevent apoptosis
	by inhibiting caspases (8).

Gene ID:	1409
NCBI Accession:	NP_000385
UniProt:	P02489
Pathways:	M Phase

## Application Details

Application Notes:	<ul> <li>WB (1:2000)</li> <li>ICC/IF (1:100)</li> <li>optimal dilutions for assays should be determined by the user.</li> </ul>
Comment:	0.5 μg/ml was sufficient for detection of 100 ng purified alphaA crystalline by colorimetric immunoblot analysis using Goat Anti-Mouse IgG:HRP as the secondary.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.2, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated


Preservative: Sodium azide

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN2481837 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

i II	•
lond	lina
land	
ana	mg

Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Conjugated antibodies should be stored at 4°C

### Images




#### Immunofluorescence (fixed cells)

**Image** 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Alpha A Crystallin Monoclonal Antibody, Clone 1H3.B8 . Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-Alpha A Crystallin Monoclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60min RT, 5min RT. Localization: Cytoplasm . Magnification: 60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) Alpha A Crystallin Antibody (D) Composite.

#### Western Blotting

**Image 2.** Western Blot analysis of Bovine tissue lysate showing detection of Alpha A Crystallin protein using Mouse Anti-Alpha A Crystallin Monoclonal Antibody, Clone 1H3.B8 . Load: 15 µg. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-Alpha A Crystallin Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT. This blot shows absolute specificity as Left: Alpha A Crystallin, Right: Alpha B Crystallin.



Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN2481837 | 09/11/2023 | Copyright antibodies-online. All rights reserved.