

Datasheet for ABIN2481896  
**anti-DNAJB1 antibody (APC)**



[Go to Product page](#)

2 Images

## Overview

Quantity:	100 µg
Target:	DNAJB1
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This DNAJB1 antibody is conjugated to APC
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunofluorescence (IF), Immunocytochemistry (ICC)

## Product Details

Immunogen:	Recombinant Protein HSP40 (Hdj1)
Clone:	3B9-E6
Isotype:	IgG1
Specificity:	Detects ~40 kDa. Does not cross-react with HDJ2 or YDJ1.
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Protein G Purified

## Target Details

Target:	DNAJB1
Alternative Name:	HSP40 ( <a href="#">DNAJB1 Products</a> )

## Target Details

---

**Background:** Human HSP40/DnaJ proteins comprise a large protein family, members of which feature the J domain (named after the bacterial DnaJ protein) (1). The J-domain spans the first 75 N-terminal amino acids and is separated from the C-terminal by a glycine/phenylalanine-rich domain (2). Members of the HSP40/DnaJ family play diverse roles in many cellular processes, such as folding, translocation, degradation and assembly of multi-protein complexes. In particular, Hdj1, the first human HSP40/DnaJ protein identified, plays an important role in protein translation and folding, as well as in the regulation of HSP70 function (3). HSP40 stimulates the ATPase activity of HSP70 which in turn causes conformational changes of the unfolded proteins (4, 5). The HSP40-HSP70-unfolded protein complex further binds to co-chaperones Hip, Hop and HSP90 which leads to protein folding, or components of protein degradation machinery CHIP and BAG-1 (6). Some studies have shown that the difference between HDJ1 and type 1 DNAJ proteins including HDJ2 and yeast Ydj1 is the result of the possession of a zinc finger domain by the latter, which helps in the function of protein folding. (7, 8).

---

**Gene ID:** 3337

---

**NCBI Accession:** [NP\\_006136](#)

---

**UniProt:** [P25685](#)

## Application Details

---

**Application Notes:**

- WB (1:2000)
- ICC/IF (1:100)
- optimal dilutions for assays should be determined by the user.

---

**Comment:** 0.5 µg/ml of ABIN2481896 was sufficient for detection of HSP40 (HDJ1) in 15 µg of HeLa cell lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.

---

**Restrictions:** For Research Use only

## Handling

---

**Format:** Liquid

---

**Concentration:** 1 mg/mL

---

**Buffer:** PBS pH 7.2, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated

---

**Preservative:** Sodium azide

---


## Handling

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C


Storage Comment: Conjugated antibodies should be stored at 4°C

## Images



### Immunofluorescence (fixed cells)

**Image 1.** Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Hsp40 Monoclonal Antibody, Clone 3B9.E6 . Tissue: HaCaT cells. Species: Human. Fixation: Cold 100% methanol for 10 minutes at -20°C. Primary Antibody: Mouse Anti-Hsp40 Monoclonal Antibody at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT.



### Western Blotting

**Image 2.** Western Blot analysis of Human Cell lysates showing detection of Hsp40 protein using Mouse Anti-Hsp40 Monoclonal Antibody, Clone 3B9.E6 . Load: 15 μg. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-Hsp40 Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.