
antibodies -online.com





anti-HSPA4 antibody (Atto 488)

2 Images



Go to Product page

Overview

Quantity:	100 μg
Target:	HSPA4
Reactivity:	Hamster
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HSPA4 antibody is conjugated to Atto 488
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	Synthetic peptide derived from the sequence of hamster HSP110, sequence identical to human and mouse
Specificity:	Detects ~110 kDa.
Cross-Reactivity:	Cow, Hamster, Human, Monkey, Mouse, Rat, Saccharomyces cerevisiae, Shark, Sheep
Purification:	Protein A Purified

Target Details

Target:	HSPA4
Alternative Name:	HSP110 (HSPA4 Products)
Background:	HSP110 belongs to a family of large stress proteins known as the HSP110/SSE Family. The
	proteins in this family are the most distantly known relatives of the well studied HSP70 family.

oinding domain (1).
osol (2). In mammals,
in the brain. Both HSP70
onstrate that the
y be mediated by an
hat HSP110 can be used


Gene ID:	10808
NCBI Accession:	NP_006635
UniProt:	Q92599

Application Details

Application Notes:	 WB (1:1000) IHC (1:100) optimal dilutions for assays should be determined by the user.
Comment:	1 μg/ml of ABIN2481905 was sufficient for detection of HSP110 in 10 μg of human cell line mixed lysate by colorimetric immunoblot analysis using Goat anti-rabbit IgG:HRP as the
	secondary antibody.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.2, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Conjugated antibodies should be stored at 4°C



Immunohistochemistry

Image 1. Immunohistochemistry analysis using Rabbit Anti-HSP110 Polyclonal Antibody . Tissue: backskin. Species: Mouse. Fixation: Bouin's Fixative Solution. Primary Antibody: Rabbit Anti-HSP110 Polyclonal Antibody at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Rabbit (green) at 1:50 for 1 hour at RT. Localization: Positive epidermal, dermal and muscle staining.



Western Blotting

Image 2. Western blot analysis of Human Cell line lysates showing detection of HSP110 protein using Rabbit Anti-HSP110 Polyclonal Antibody . Load: 15 μg protein. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Rabbit Anti-HSP110 Polyclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Donkey Anti-Rabbit IgG: HRP for 1 hour at RT.