

Datasheet for ABIN2481921

anti-HSPE1 antibody (AA 91-101) (Atto 390)

2 Images



Go to Product page

| _ | | | | | | |
|---|-----|----------|----|----|----|----|
| | 1// | Д | rv | 16 | ٦/ | ٨ |
| U | W | \vdash | ΙV | Ιt | ٦, | /V |

| Quantity: | 100 μg |
|----------------------|--|
| Target: | HSPE1 |
| Binding Specificity: | AA 91-101 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This HSPE1 antibody is conjugated to Atto 390 |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunofluorescence (IF), Immunocytochemistry (ICC) |

Product Details

| Immunogen: | Human Cpn10 peptide AA 91-101 | |
|-------------------|---|--|
| Specificity: | Detects ~ 10 kDa. | |
| Cross-Reactivity: | Cow, Dog, Guinea Pig, Human, Mouse, Pig, Rabbit, Rat, Sheep, Xenopus laevis | |
| Purification: | Protein A Purified | |

Target Details

| Target: | HSPE1 |
|-------------------|--|
| Alternative Name: | Cpn10 (HSPE1 Products) |
| Background: | Chaperonin 10, otherwise known as Cpn10, (groES in E.coli) make up a family of small heart |

| situations such as pregnancy (4). |
|---|
| development. This identification suggest that Cpn10 may act like a hormone in stressful |
| identical with early pregnancy factor, which is involved in control over cell growth and |
| also present in human maternal serum (2, 3). It has been reported that human Cpn10 is |
| protein folding and assembly process (1). Cpn10 has been located in human platelets, but is |
| Cpn10 and Cpn60 both exhibit sevenfold axis of symmetry and function as a team in the |
| chaperone and interacts with the HSP60 family to promote proper folding of polypeptides. |
| shock proteins with an approximate molecular mass of 10 kDa (HSP10s). Cpn10 acts as a co- |

Gene ID: 3336

NCBI Accession: NP_002148

UniProt: P61604

Pathways: Positive Regulation of Endopeptidase Activity

Application Details

Application Notes: • WB (1:1000)

• IHC (1:100)

• optimal dilutions for assays should be determined by the user.


Comment: $1~\mu\text{g/ml of ABIN2481921 was sufficient for detection of Cpn10 in 10~\mu\text{g of Hela lysates by}}$

colorimetric immunoblot analysis using Goat anti-rabbit IgG:HRP as the secondary antibody.

Restrictions: For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 mg/mL |
| Buffer: | PBS pH 7.3, 0.02 % sodium azide, Storage buffer may change when conjugated |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C |
| Storage Comment: | Conjugated antibodies should be stored at 4°C |





Immunohistochemistry

Image 1. Immunohistochemistry analysis using Rabbit Anti-Cpn10 Polyclonal Antibody . Tissue: backskin. Species: Mouse. Fixation: Bouin's Fixative Solution. Primary Antibody: Rabbit Anti-Cpn10 Polyclonal Antibody at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Rabbit (green) at 1:50 for 1 hour at RT. Localization: Strong basal layer punctate staining in the cytoplasm, weakens as it ascends to upper epidermis..

Western Blotting

Image 2. Western blot analysis of Rat brain cell lysates showing detection of Cpn10 protein using Rabbit Anti-Cpn10 Polyclonal Antibody . Load: 15 µg protein. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Rabbit Anti-Cpn10 Polyclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Donkey Anti-Rabbit IgG: HRP for 1 hour at RT.