
# antibodies - online.com







# anti-SLC38A1 antibody (AA 1-63) (Biotin)

**Images** 



# Overview

Quantity:	100 μg
Target:	SLC38A1
Binding Specificity:	AA 1-63
Reactivity:	Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This SLC38A1 antibody is conjugated to Biotin
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)

# **Product Details**

Immunogen:	Fusion protein amino acids 1- 63 of rat SNAT1
Clone:	S104-32
Isotype:	lgG1
Specificity:	Detects ~50 kDa.
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Protein G Purified


# **Target Details**


Target: SLC38A1
-----------------

# **Target Details**

Target Details	
Alternative Name:	SLC38A1 (SLC38A1 Products)
Background:	The sodium-coupled neutral amino acid transporters (SNAT) of the SLC38 gene family include
	System A subtypes SNAT1, SNAT2 and SNAT4 and System N subtypes SNAT3 and SNAT5.
	The SLC38 transporters are essential for the uptake of nutrients, energy production,
	metabolism, detoxification, and the cycling of neurotransmitters. The SNAT1 protein, also
	designated ATA1 or NAT2 is encoded by the human gene SLC38A1 which maps to
	chromosome 12q13.11. SNAT1 is responsible for the transport of glutamine, an intermediate in
	the synthesis of urea, and may be involved in the generation of glutamate in the retina. SNAT1
	protein may be detected in some tissues such as heart, brain and placenta and expression
	levels are enriched in certain neuronal populations within the CNS. SNAT1 is not present in
	astrocytes.
Gene ID:	170567
NCBI Accession:	NP_620187
UniProt:	Q9JM15
Application Details	
Application Notes:	• WB (1:1000)
	optimal dilutions for assays should be determined by the user.
Comment:	1 μg/ml of ABIN2482519 was sufficient for detection of SNAT1 in 20 μg of lysates from
	neocortical neurons cultured under amino acid starvation conditions and assayed by
	colorimetric immunoblot analysis using goat anti-mouse IgG:HRP as the secondary antibody.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C

# **Images**






### **Immunocytochemistry**

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-SNAT1 Monoclonal Antibody, Clone S104-32 (ABIN2482519). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-SNAT1 Monoclonal Antibody (ABIN2482519) at 1:200 for overnight at 4 °C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) SNAT1 Antibody (D) Composite.

### Immunofluorescence (fixed cells)

Image 2. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-SNAT1 Monoclonal Antibody, Clone S104-32. Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-SNAT1 Monoclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60min RT, 5min RT. Localization: Cell Membrane. Magnification: 60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) SNAT1 Antibody (D) Composite.



# **Western Blotting**

**Image 3.** Western Blot analysis of Rat brain membrane lysate showing detection of SLC38A1 protein using Mouse Anti-SLC38A1 Monoclonal Antibody, Clone S104-32. Primary Antibody: Mouse Anti-SLC38A1 Monoclonal Antibody at 1:1000.