
antibodies - online.com







anti-SCN9A antibody (AA 1751-1946) (PE)



Images



\sim	
()\/\Di	view
	VICVV

Quantity:	100 μg
Target:	SCN9A
Binding Specificity:	AA 1751-1946
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This SCN9A antibody is conjugated to PE
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunofluorescence (IF), Immunoprecipitation (IP), Immunocytochemistry (ICC), Antibody Array (AA)

Product Details

Immunogen:	Fusion protein amino acids 1751-1946 (C-terminus) of human Nav1.7
Clone:	N68-6 (Formerly S68-6)
Isotype:	lgG1
Specificity:	Detects ~230 kDa. No cross-reactivity against other Nav channels.
Cross-Reactivity:	Hamster, Human, Mouse, Rat
Purification:	Protein G Purified

Target Details


Target Details


Alternative Name:	Nav1.7 (SCN9A Products)
Background:	Nav1.7 is a voltage-gated sodium channel and plays a critical role in the generation and
	conduction of action potentials and is thus important for electrical signaling by most excitable
	cells. Therapeutically, the association of pain insensitivity with the loss of function of a certain
	sodium channel may have implications. Since Nav1.7 is not present in cardiac muscle or
	neurons in the central nervous system, blockers of Nav1.7 will not have direct action on these
	cells and thus can have less side effects than current pain medications. By performing more
	studies, there is a possibility to develop a new generation of drugs that can reduce the pain
	intensity in animals (1-3).
Gene ID:	6335
NCBI Accession:	NP_002968
UniProt:	Q15858
Application Details	
Application Notes:	• WB (1:1000)
	• IHC (1:1000)
	• ICC/IF (1:100)
	optimal dilutions for assays should be determined by the user.
Comment:	1 μg/ml of ABIN2483102 was sufficient for detection of Nav1.7 in 10 μg of HEK-293 cell lysat
	transiently expressing Nav1.7 by colorimetric immunoblot analysis using Goat anti-mouse
	IgG:HRP as the secondary antibody.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C

Storage Comment:

Conjugated antibodies should be stored at 4°C

Images






Immunocytochemistry

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Nav1.7 Monoclonal Antibody, Clone N68/6 (ABIN2483102). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-Nav1.7 Monoclonal Antibody (ABIN2483102) at 1:100 for overnight at 4 °C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) Nav1.7 Antibody (D) Composite.

Western Blotting

Image 2. Western Blot analysis of hamster CHO cells showing detection of Nav1.7 Sodium Channel protein using Mouse Anti-Nav1.7 Sodium Channel Monoclonal Antibody, Clone S68-6 . Load: 15 μg. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-Nav1.7 Sodium Channel Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.



Immunohistochemistry

Image 3. Immunohistochemistry analysis using Mouse Anti-Nav1.7 Sodium Channel Monoclonal Antibody, Clone S68-6. Tissue: backskin. Species: Mouse. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-Nav1.7 Sodium Channel Monoclonal Antibody at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT.

Please check the product details page for more images. Overall 5 images are available for ABIN2483102.