# antibodies .- online.com





## anti-KCNQ1 antibody (AA 2-101) (PE)



## **Images**



Go to Product page

$\sim$	
( )\/\	rview
$\cup$	1 410 44

Quantity:	100 μg
Target:	KCNQ1
Binding Specificity:	AA 2-101
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This KCNQ1 antibody is conjugated to PE
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunofluorescence (IF), Immunocytochemistry (ICC), Antibody Array (AA)

## **Product Details**

**Target Details** 


KCNQ1

Target:

Immunogen:	Fusion protein amino acids 2-101 of human KCNQ1
Clone:	N37A-10 (Formerly S37A-10)
Isotype:	lgG1
Specificity:	Detects ~75 kDa.
Cross-Reactivity:	Hamster, Human, Mouse, Rat
Purification:	Protein G Purified


## **Target Details**

rai get Betaile	
Alternative Name:	KCNQ1 (KCNQ1 Products)
Background:	Kv7.1 (KvLQT1) is a potassium channel protein coded for by the gene KCNQ1. Kv7.1 is present
	in the cell membranes of cardiac muscle tissue and in inner ear neurons (1) among other
	tissues. In the cardiac cells, Kv7.1 mediates the IKs (or slow delayed rectifying K+) current that
	contributes to the repolarization of the cell, terminating the cardiac action potential and thereby
	the heart's contraction (2, 3).
Gene ID:	3784
NCBI Accession:	NP_000209
UniProt:	P51787
Pathways:	Negative Regulation of Hormone Secretion, Sensory Perception of Sound
Application Details	
Application Notes:	• WB (1:1000)
	• IHC (1:1000)
	<ul> <li>ICC/IF (1:100)</li> <li>optimal dilutions for assays should be determined by the user.</li> </ul>
	optimal unutions for assays should be determined by the user.
Comment:	1 μg/ml of ABIN2483170 was sufficient for detection of KCNQ1 in 10 μg of COS-1 cell lysate
	transiently expressing KCNQ1 by colorimetric immunoblot analysis using Goat anti-mouse
	IgG:HRP as the secondary antibody.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Conjugated antibodies should be stored at 4°C




### **Immunocytochemistry**

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-KCNQ1 Monoclonal Antibody, Clone N37A/10 (ABIN2483170). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-KCNQ1 Monoclonal Antibody (ABIN2483170) at 1:100 for overnight at 4 °C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) KCNQ1 Antibody (D) Composite.



### **Immunohistochemistry**

Image 2. Immunohistochemistry analysis using Mouse Anti-KCNQ1 Monoclonal Antibody, Clone S37A-10. Tissue: Brain Slice. Species: Mouse. Fixation: 10% Formalin Solution for 12-24 hours at RT. Primary Antibody: Mouse Anti-KCNQ1 Monoclonal Antibody at 1:1000 for 1 hour at RT. Secondary Antibody: HRP/DAB Detection System: Biotinylated Goat Anti-Mouse, Streptavidin Peroxidase, DAB Chromogen (brown) for 30 minutes at RT. Counterstain: Mayer Hematoxylin (purple/blue) nuclear stain at 250-500 µl for 5 minutes at RT.



#### **Immunohistochemistry**

Image 3. Immunohistochemistry analysis using Mouse Anti-KCNQ1 Monoclonal Antibody, Clone S37A-10. Tissue: hippocampus. Species: Human. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-KCNQ1 Monoclonal Antibody at 1:1000 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT.

)	
	Please check the product details page for more images. Overall 5 images are available for ABIN2483170.