antibodies.com

Datasheet for ABIN2483200 anti-Notch1 antibody (AA 20-216) (FITC)







Overview

Quantity:	100 µg
Target:	Notch1 (NOTCH1)
Binding Specificity:	AA 20-216
Reactivity:	Mouse
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Notch1 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	Fusion protein amino acids 20-216 (extracellular N-terminus, EGF-like domains 1-5) of mouse
	Notch1
Clone:	S253-32
laaturaa	
Isotype:	lgG1
Specificity:	Detects >270 kDa. Does not cross-react with Notch 2 or Notch3.
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Protein G Purified
r unneation.	
Target Details	
Target:	Notch1 (NOTCH1)
5	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN2483200 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
Alternative Name:	Notch1 (NOTCH1 Products)
Background:	Notch is synthesized in the endoplasmic reticulum as an inactive form which is proteolytically cleaved by a furin-like convertase (S1 cleavage) in the trans-golgi network before it reaches the plasma membrane to yield an active, ligand-accessible form. Cleavage results in a C-terminal fragment N(TM) and a N-terminal fragment N(EC). Following ligand binding, it is cleaved (S2 cleavage) by TNF-alpha converting enzyme (TACE) to yield a membrane-associated intermediate fragment called Notch extracellular truncation (NEXT). This fragment is then cleaved by presenilin-dependent gamma-secretase (S3 cleavage) to release the intracellular domain (NICD) from the membrane.
Gene ID:	18128
NCBI Accession:	NP_032740
UniProt:	Q01705
Pathways:	Notch Signaling, Stem Cell Maintenance, Regulation of Muscle Cell Differentiation, Tube Formation, Skeletal Muscle Fiber Development
Application Details	
Application Notes:	WB (1:1000)optimal dilutions for assays should be determined by the user.

Comment:	1 $\mu g/ml$ of ABIN2483200 was sufficient for detection of Notch1 in 20 μg of rat brain membrane
	lysate and assayed by colorimetric immunoblot analysis using goat anti-mouse IgG:HRP as the
	secondary antibody.

Restrictions:


For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN2483200 | 09/11/2023 | Copyright antibodies-online. All rights reserved. Storage Comment:

Images






Immunocytochemistry

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Notch1 Monoclonal Antibody, Clone S253-32 (ABIN2483200). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-Notch1 Monoclonal Antibody (ABIN2483200) at 1:50 for overnight at 4 °C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) Notch1 Antibody (D) Composite.

Western Blotting

Image 2. Western Blot analysis of Rat Brain Membrane showing detection of ~270 kDa Notch1 protein using Mouse Anti-Notch1 Monoclonal Antibody, Clone S253-32 . Lane 1: MW Ladder. Lane 2: Rat Brain Membrane (10 μg). . Load: 10 μg. Block: 5% milk. Primary Antibody: Mouse Anti-Notch1 Monoclonal Antibody at 1:1000 for 1 hour at RT. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:200 for 1 hour at RT. Color Development: TMB solution for 10 min at RT. Predicted/Observed Size: ~270 kDa.



Immunofluorescence (fixed cells)

Image 3. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Notch1 Monoclonal Antibody, Clone S253-32 . Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-Notch1 Monoclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60min RT, 5min RT. Localization: Cell Membrane, Nucleus. Magnification: 60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) Notch1 Antibody (D) Composite.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 4/4 | Product datasheet for ABIN2483200 | 09/11/2023 | Copyright antibodies-online. All rights reserved.