# antibodies .- online.com







## anti-CACNA1G antibody (AA 2052-2172) (APC)

**Images** 



## Overview

| Quantity:            | 100 μg  |
|----------------------|---|
| Target:              | CACNA1G   |
| Binding Specificity: | AA 2052-2172  |
| Reactivity:          | Mouse   |
| Host:                | Mouse   |
| Clonality:           | Monoclonal  |
| Conjugate:           | This CACNA1G antibody is conjugated to APC  |
| Application:         | Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC) |

## **Product Details**

| Immunogen:        | Fusion protein amino acids 2052-2172 (cytoplasmic C-terminus) of mouse Cav3.1 |
|-------------------|---|
| Clone:            | S178A-9   |
| Isotype:          | lgG1  |
| Specificity:      | Detects ~<200 kDa. Does not cross-react with Cav3.2.                          |
| Cross-Reactivity: | Human, Mouse, Rat   |
| Purification:     | Protein G Purified  |

## **Target Details**

Target: CACNA1G


## **Target Details**


| Alternative Name:   | Cav3.1 (CACNA1G Products)   |
|---------------------|---|
| Background:         | Calcium channel CaV3.1 (a1G) is a low-voltage-activated T-type calcium channel. Such T-type channels are expressed throughout the body. In the heart, they may be involved in pacemaker current. In neurons, these channels may play a secondary pacemaker role (1). With the |
|                     | ubiquitous expression, it is not surprising that alterations in channel function have been  |
|                     | implicated in disease. Drugs that act to block T-type calcium channels are used as anti-  |
|                     | hypertensives, antiepileptic's, and blocking of T-type calcium channels may be involved in the  |
|                     | action of some anesthetics and antipsychotics as well (1). Much remains to be determined  |
|                     | about the precise cellular localization, in vivo physiological roles, roles in disease states and   |
|                     | possible routes to modulate their structure/function to ameliorate effects of disease.  |
| Gene ID:            | 12291   |
| NCBI Accession:     | NP_001106284  |
| UniProt:            | Q9WUT2  |
| Application Details |   |
| Application Notes:  | <ul><li>WB (1:1000)</li><li>ICC/IF (1:100)</li></ul>  |
|                     | optimal dilutions for assays should be determined by the user.  |
| Comment:            | 1 μg/ml of ABIN2483300 was sufficient for detection of Cav3.1 in 20 μg of rat brain membrane  |
|                     | lysate and assayed by colorimetric immunoblot analysis using goat anti-mouse IgG:HRP as the   |
|                     | secondary antibody.   |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Format:             | Liquid  |
| Concentration:      | 1 mg/mL   |
| Buffer:             | PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated   |
| Preservative:       | Sodium azide  |
| Precaution of Use:  | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.  |
| Storage:            | 4 °C  |
|                     |   |

Storage Comment:

Conjugated antibodies should be stored at 4°C

#### **Images**






#### **Immunocytochemistry**

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Cav3.1 Monoclonal Antibody, Clone S178A-9 (ABIN2483300). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-Cav3.1 Monoclonal Antibody (ABIN2483300) at 1:50 for overnight at 4 °C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) Cav3.1 Antibody (D) Composite.

## Western Blotting

**Image 2.** Western Blot analysis of Rat brain membrane lysate showing detection of Cav3.1 Calcium Channel protein using Mouse Anti-Cav3.1 Calcium Channel Monoclonal Antibody, Clone S178A-9. Primary Antibody: Mouse Anti-Cav3.1 Calcium Channel Monoclonal Antibody at 1:1000.



## Immunofluorescence (fixed cells)

Image 3. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Cav3.1 Monoclonal Antibody, Clone S178A-9. Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-Cav3.1 Monoclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:200 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60 min at RT, 5 min at RT. Localization: Cell Membrane, Membrane, Cytoplasm, Nucleoplasm. Magnification: 60X.