

Datasheet for ABIN2483697

anti-ASIC1 antibody (AA 460-526) (PE)**3** Images[Go to Product page](#)

Overview

| | |
|----------------------|---|
| Quantity: | 100 µg |
| Target: | ASIC1 (ACCN2) |
| Binding Specificity: | AA 460-526 |
| Reactivity: | Mouse |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This ASIC1 antibody is conjugated to PE |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF) |

Product Details

| | |
|-------------------|--|
| Immunogen: | Fusion protein amino acids 460-526 (Cytoplasmic C-terminus) of mouse ASIC1 |
| Clone: | S271-44 |
| Isotype: | IgG1 |
| Specificity: | Detects ~60 kDa. |
| Cross-Reactivity: | Human, Mouse, Rat |
| Purification: | Protein G Purified |

Target Details

| | |
|---------|---------------|
| Target: | ASIC1 (ACCN2) |
|---------|---------------|

Target Details


| | |
|-------------------|--|
| Alternative Name: | ASIC1 (ACCN2 Products) |
| Background: | Acid sensing ion channel ASIC1, also designated ACCN2, BNAC2 and ASIC1a, is present in brain as a 4.3-kb transcript with localization to rat dorsal root ganglia. Insitu hybridization of rat brain suggests that ASIC1 is most abundant in the main olfactory bulb, cerebral cortex, hippocampal formation, habenula, basolateral amygdaloid nuclei and cerebellum. ASIC1 and H ⁺ -gated currents may contribute to the development of fear and anxiety. |
| Gene ID: | 11419 |
| NCBI Accession: | NP_033727 |
| UniProt: | Q6NXK8 |

Application Details

| | |
|--------------------|--|
| Application Notes: | <ul style="list-style-type: none">• WB (1:1000)• optimal dilutions for assays should be determined by the user. |
| Comment: | 1 µg/ml of ABIN2483697 was sufficient for detection of ASIC1 in 20 µg of rat brain lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody. |
| Restrictions: | For Research Use only |


Handling

| | |
|--------------------|--|
| Format: | Liquid |
| Concentration: | 1 mg/mL |
| Buffer: | PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C |
| Storage Comment: | Conjugated antibodies should be stored at 4°C |




Immunocytochemistry

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-ASIC1 Monoclonal Antibody, Clone S271-44 (ABIN2483697). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-ASIC1 Monoclonal Antibody (ABIN2483697) at 1:100 for overnight at 4 °C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) ASIC1 Antibody (D) Composite.



Western Blotting

Image 2. Western Blot analysis of Rat brain lysates showing detection of ASIC1 protein using Mouse Anti-ASIC1 Monoclonal Antibody, Clone S271-44 . Primary Antibody: Mouse Anti-ASIC1 Monoclonal Antibody at 1:100, 1:250, 1:500, and 1:1000.



Immunofluorescence (fixed cells)

Image 3. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-ASIC1 Monoclonal Antibody, Clone S271-44 . Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-ASIC1 Monoclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60min RT, 5min RT. Localization: Cell Membrane, Nucleus. Magnification: 60X.

(A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) ASIC1 Antibody (D) Composite.