

Datasheet for ABIN2483744

anti-Solute Carrier Family 17 (Vesicular Glutamate Transporter), Member 6 (SLC17A6) (AA 501-582) antibody (FITC)



[Go to Product page](#)

3 Images

Overview

Quantity:	100 µg
Target:	Solute Carrier Family 17 (Vesicular Glutamate Transporter), Member 6 (SLC17A6)
Binding Specificity:	AA 501-582
Reactivity:	Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	FITC
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	Fusion protein amino acids 501-582 (cytoplasmic C-terminus) of rat VGlut2
Clone:	S29-29
Isotype:	IgG1
Specificity:	Detects ~60 kDa.
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Protein G Purified

Target Details

Target:	Solute Carrier Family 17 (Vesicular Glutamate Transporter), Member 6 (SLC17A6)
Alternative Name:	VGLUT2 (SLC17A6 Products)
Background:	<p>The ATP-dependent, chloride-sensitive vesicular glutamate transporters (VGLUT) include BNPI (VGLUT1), VGLUT2 (DNPI) and VGLUT3. The brain expresses BNPI (brain specific Na⁺-dependent inorganic phosphate (Pi) cotransporter) and VGLUT2 in a complementary fashion. The telencephalic regions express BNPI, whereas the lower brainstem and diencephalic regions express VGLUT2. Rat pinealocytes express both BNPI and VGLUT2. The striatum, hippocampus, cerebral cortex and raphe nuclei express VGLUT3 in a small number of neurons. Pancreatic α and β cells express BNPI and VGLUT2 in response to glucose concentrations. Human VGLUT3 shares a 72 % sequence homology with VGLUT2 and BNPI.</p>
Gene ID:	84487
NCBI Accession:	NP_445879
UniProt:	Q9J112

Application Details


Application Notes:	<ul style="list-style-type: none">• WB (1:1000)• ICC/IF (1:100)• optimal dilutions for assays should be determined by the user.
Comment:	1 μ g/ml of ABIN2483744 was sufficient for detection of VGLut2 in 20 μ g of rat brain lysate by colorimetric immunoblot analysis using goat anti-mouse IgG:HRP as the secondary antibody.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C


Storage Comment: Conjugated antibodies should be stored at 4°C

Validation report #103875 for Immunofluorescence (IF)




Immunocytochemistry

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-VGLUT2 Monoclonal Antibody, Clone S29-29 (ABIN2483744). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-VGLUT2 Monoclonal Antibody (ABIN2483744) at 1:50 for overnight at 4 °C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) VGLUT2 Antibody (D) Composite.



Immunofluorescence (fixed cells)

Image 2. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-VGLUT2 Monoclonal Antibody, Clone S29-29 . Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-VGLUT2 Monoclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:200 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60 min at RT, 5 min at RT. Localization: Cytoplasmic Vesicle, Secretory Vesicle, Synaptic Vesicle Membrane, Membrane, Cell Junction, Synapse. Magnification: 60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) VGLUT2 Antibody (D) Composite.



Western Blotting

Image 3. Western Blot analysis of Rat brain membrane lysate showing detection of VGLUT2 protein using Mouse Anti-VGLUT2 Monoclonal Antibody, Clone S29-29 . Primary Antibody: Mouse Anti-VGLUT2 Monoclonal Antibody at 1:000.