antibodies

Datasheet for ABIN2483810 anti-AMIGO1 antibody (AA 554-574) (APC)

3 Images



Overview

Quantity:	100 µg
Target:	AMIG01
Binding Specificity:	AA 554-574
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This AMIGO1 antibody is conjugated to APC
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details


Immunogen:	Fusion protein amino acids 554-574 (cytoplasmic C-terminus) of human AMIGO-1
Clone:	S86-36
Isotype:	lgG1
Specificity:	Detects ~60-80 kDa depending on maturity/glycosylation.
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Protein G Purified
Target Details	
Target:	AMIG01

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN2483810 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
Alternative Name:	AMIG01 (AMIG01 Products)
Background:	The amphoterin-induced gene and ORF (AMIGO) family of proteins consists of AMIGO1, AMIGO2 and AMIGO3. All three members are single pass type I membrane proteins that contain several leucine-rich repeats, one IgG domain and a transmembrane domain. The AMIGO proteins are specifically expressed on fiber tracts of neuronal tissues and participate in their formation. They can form complexes with each other, but can also self-bind. AMIGO1, also designated Alivin2, promotes growth and fasciculation of neurites and plays a role in myelination and fasciculation of developing neural axons. In cerebellar neurons, AMIGO2 (Alivin1) is crucial for depolarization-dependent survival. Similar to AMIGO1 and AMIGO2, AMIGO3 (Alivin3) plays a role in hemophilic and/or heterophilic cell-cell interaction and signal transduction.
Gene ID:	57463
NCBI Accession:	NP_065754
UniProt:	Q86WK6
Application Details	
Application Notes:	WB (1:1000)optimal dilutions for assays should be determined by the user.
Comment:	1 μg/ml of ABIN2483810 was sufficient for detection of AMIGO-1 in 20 μg of rat brain membrane lysate and assayed by colorimetric immunoblot analysis using goat anti-mouse IgG:HRP as the secondary antibody.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN2483810 | 09/11/2023 | Copyright antibodies-online. All rights reserved. Storage Comment:

Images



kDa MW 2 150 100 75 − 50 − 2 4 ~60-80 kDa AMIGO-1


Immunocytochemistry

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-AMIGO-1 Monoclonal Antibody, Clone S86-36 (ABIN2483810). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-AMIGO-1 Monoclonal Antibody (ABIN2483810) at 1:200 for overnight at 4 °C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) AMIGO-1 Antibody (D) Composite.

Western Blotting

Image 2. Western Blot analysis of Rat Brain Membrane showing detection of 60-80 kDa AMIGO-1 protein using Mouse Anti-AMIGO-1 Monoclonal Antibody, Clone S86-36 . Lane 1: Molecular Weight Ladder. Lane 2: Rat Brain Membrane. Load: 15 µg. Block: 2% BSA and 2% Skim Milk in 1X TBST. Primary Antibody: Mouse Anti-AMIGO-1 Monoclonal Antibody at 1:200 for 16 hours at 4°C. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:1000 for 1 hour RT. Color Development: ECL solution for 6 min in RT. Predicted/Observed Size: 60-80 kDa.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/4 | Product datasheet for ABIN2483810 | 09/11/2023 | Copyright antibodies-online. All rights reserved.



Immunofluorescence (fixed cells)

Image 3. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-AMIGO-1 Monoclonal Antibody, Clone S86-36 . Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-AMIGO-1 Monoclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60min RT, 5min RT. Localization: Cell Membrane, Nucleus. Magnification: 60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) AMIGO-1 Antibody (D) Composite.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 4/4 | Product datasheet for ABIN2483810 | 09/11/2023 | Copyright antibodies-online. All rights reserved.