
antibodies -online.com







anti-HCN1 antibody (AA 778-910) (Atto 390)





Go to Product page

\sim						
	1//	Д	r۱	1	Θ 1	٨

Quantity:	100 μg	
Target:	HCN1	
Binding Specificity:	AA 778-910	
Reactivity:	Rat	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This HCN1 antibody is conjugated to Atto 390	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Antibody Array (AA)	

Product Details

Immunogen:	Fusion protein amino acids 778-910 (C terminus) of rat HCN1	
Clone:	S70	
Isotype:	lgG1	
Specificity:	Detects ~100 kDa. No cross-reactivity against HCN2.	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Protein G Purified	

Target Details

	Target:	HCN1
--	---------	------

Target Details


rarget Details		
Alternative Name:	HCN1 (HCN1 Products)	
Background:	Hyperpolarization-activated cation channels of the HCN gene family, such as HCN1, play a crucial role in the regulatons of cell excitability. Importantly, they contribute to spontaneous rhythmic activity in both the heart and brain (1).	
Gene ID:	84390	
NCBI Accession:	NP_445827	
UniProt:	Q9JKB0	
Pathways:	Asymmetric Protein Localization	
Application Details		
Application Notes:	 WB (1:1000) IHC (1:1000) ICC/IF (1:100) optimal dilutions for assays should be determined by the user. 	
Comment:	1 μg/ml of ABIN2483954 was sufficient for detection of HCN1 in 10 μg of rat brain lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated	
Preservative:	Sodium azide	
Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTAN should be handled by trained staff only.		

Conjugated antibodies should be stored at 4°C

4°C


Storage:

Storage Comment:




Immunohistochemistry

Image 1. Immunohistochemistry analysis using Mouse Anti-HCN1 Monoclonal Antibody, Clone S70-28. Tissue: Cerebellum. Species: Mouse. Fixation: 10% Formalin Solution for 12-24 hours at RT. Primary Antibody: Mouse Anti-HCN1 Monoclonal Antibody at 1:1000 for 1 hour at RT. Secondary Antibody: HRP/DAB Detection System: Biotinylated Goat Anti-Mouse, Streptavidin Peroxidase, DAB Chromogen (brown) for 30 minutes at RT. Counterstain: Mayer Hematoxylin (purple/blue) nuclear stain at 250-500 µl for 5 minutes at RT. Localization: Cytoplasmic staining of Purkinje cells.



Western Blotting

Image 2. Western Blot analysis of Rat brain membrane lysate showing detection of HCN1 protein using Mouse Anti-HCN1 Monoclonal Antibody, Clone S70-28 . Load: 15 μg. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-HCN1 Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.



Immunohistochemistry

Image 3. Immunohistochemistry analysis using Mouse Anti-HCN1 Monoclonal Antibody, Clone S70-28. Tissue: hippocampus. Species: Human. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-HCN1 Monoclonal Antibody at 1:1000 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT.

Please check the product details page for more images. Overall 5 images are available for ABIN2483954.